





VWF Antibody, FITC conjugated

Product Code	CSB-PA12779C0Rb
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P04275
Immunogen	Recombinant Human von Willebrand factor protein (1-273AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	Important in the maintenance of hemostasis, it promotes adhesion of platelets to the sites of vascular injury by forming a molecular bridge between subendothelial collagen matrix and platelet-surface receptor complex GPIb-IX-V. Also acts as a chaperone for coagulation factor VIII, delivering it to the site of injury, stabilizing its heterodimeric structure and protecting it from premature clearance from plasma.
Form	Liquid
Form Conjugate	Liquid FITC
	·
Conjugate	FITC Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Conjugate Storage Buffer Purification Method Isotype	FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal von Willebrand factor (vWF) [Cleaved into: von Willebrand antigen 2 (von
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal von Willebrand factor (vWF) [Cleaved into: von Willebrand antigen 2 (von Willebrand antigen II)], VWF, F8VWF