





## VTCN1 Antibody, FITC conjugated

Product Code	CSB-PA13005C0Rb
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q7Z7D3
Immunogen	Recombinant Human V-set domain-containing T-cell activation inhibitor 1 protein (26-258AA)
Raised In	Rabbit
Species Reactivity	Human
<b>Tested Applications</b>	ELISA
Relevance	Negatively regulates T-cell-mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. When expressed on the cell surface of tumor macrophages, plays an important role, together with regulatory T-cells (Treg), in the suppression of tumorassociated antigen-specific T-cell immunity. Involved in promoting epithelial cell transformation.
Form	Liquid
Form Conjugate	Liquid FITC
	· · · · · · · · · · · · · · · · · · ·
Conjugate	FITC Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	FITC  Preservative: 0.03% Proclin 300  Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	FITC  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified
Conjugate Storage Buffer Purification Method Isotype	FITC  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified  IgG
Conjugate Storage Buffer  Purification Method Isotype Clonality	FITC  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified  IgG  Polyclonal  V-set domain-containing T-cell activation inhibitor 1 (B7 homolog 4) (B7-H4) (B7h.5) (Immune costimulatory protein B7-H4) (Protein B7S1) (T-cell
Conjugate Storage Buffer  Purification Method Isotype Clonality Alias	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified  IgG  Polyclonal  V-set domain-containing T-cell activation inhibitor 1 (B7 homolog 4) (B7-H4) (B7h.5) (Immune costimulatory protein B7-H4) (Protein B7S1) (T-cell costimulatory molecule B7x), VTCN1, B7H4