






Phospho-BAD (Ser91/128) Antibody

Uniprot No. Q92934 Immunogen Peptide sequence around phosphorylation site of Serine 91 (E-P-S(p)-P-F) derived from Human BAD or Serine 128 (E-P-S(p)-P-F) derived from Mouse BAD. Raised In Rabbit Rabbit Specificity The antibody detects endogenous levels of BAD only when phosphorylated at serine 91/ serine 128. Tested Applications ELISA,WB,IHC;WB:1:500-1:1000,IHC:1:50-1:100 Relevance The protein encoded by BAD gene is a member of the BCL-2 family members are known to be regulators of programmed cell death. This protein postively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calciment were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. Wang HG., Cell 87:629-638(1996). Ottlile S., J. Biol. Chem. 272:30866-30872(1997). The MGC Project Team, Genome Res. 14:2121-2127(2004). Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl., 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter; Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names		
Uniprot No. Q92934	Product Code	CSB-PA148354
Peptide sequence around phosphorylation site of Serine 91 (E-P-S(p)-P-F) derived from Human BAD or Serine 128 (E-P-S(p)-P-F) derived from Mouse BAD.	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
derived from Human BAD or Serine 128 (E-P-S(p)-P-F) derived from Mouse BAD. Raised In Rabbit Species Reactivity Human, Mouse, Rat Specificity The antibody detects endogenous levels of BAD only when phosphorylated at serine 91/ serine 128. Tested Applications ELISA, WB, IHC, WB:1:500-1:1000, IHC:1:50-1:100 Relevance The protein encoded by BAD gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. Wang HG., Cell 87:629-638(1996). Ottilie S., J. Biol. Chem. 272:30866-30872(1997). The MGC Project Team, Genome Res. 14:2121-2127(2004). Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter; Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names	Uniprot No.	Q92934
Specificity Human,Mouse,Rat Specificity The antibody detects endogenous levels of BAD only when phosphorylated at serine 91/ serine 128. Tested Applications ELISA,WB,IHC;WB:1:500-1:1000,IHC:1:50-1:100 Relevance The protein encoded by BAD gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. Wang HG., Cell 87:629-638(1996). Ottilie S., J. Biol. Chem. 272:30866-30872(1997). The MGC Project Team, Genome Res. 14:2121-2127(2004). Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter; Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names	Immunogen	derived from Human BAD or Serine 128 (E-P-S(p)-P-F) derived from Mouse
Specificity The antibody detects endogenous levels of BAD only when phosphorylated at serine 91/ serine 128. Tested Applications ELISA,WB,IHC;WB:1:500-1:1000,IHC:1:50-1:100 Relevance The protein encoded by BAD gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. Wang HG., Cell 87:629-638(1996). Ottilie S., J. Biol. Chem. 272:30866-30872(1997). The MGC Project Team, Genome Res. 14:2121-2127(2004). Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter; Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names	Raised In	Rabbit
Tested Applications ELISA,WB,IHC;WB:1:500-1:1000,IHC:1:50-1:100 Relevance The protein encoded by BAD gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. Wang HG., Cell 87:629-638(1996). Ottilie S., J. Biol. Chem. 272:30866-30872(1997). The MGC Project Team, Genome Res. 14:2121-2127(2004). Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter; Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names	Species Reactivity	Human, Mouse, Rat
Relevance The protein encoded by BAD gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. Wang HG., Cell 87:629-638(1996). Ottilie S., J. Biol. Chem. 272:30866-30872(1997). The MGC Project Team, Genome Res. 14:2121-2127(2004). Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter; Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names	Specificity	
The protein encoded by BAD gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. Wang HG., Cell 87:629-638(1996). Ottilie S., J. Biol. Chem. 272:30866-30872(1997). The MGC Project Team, Genome Res. 14:2121-2127(2004). Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter; Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names	Tested Applications	ELISA,WB,IHC;WB:1:500-1:1000,IHC:1:50-1:100
Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter; Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names BAD	Relevance	family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. Wang HG., Cell 87:629-638(1996). Ottilie S., J. Biol. Chem. 272:30866-30872(1997).
and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter; Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names BAD	Form	
Alias BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter; Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names BAD	Purification Method	and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were
Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names BAD	Clonality	Polyclonal
Species Homo sapiens (Human) Target Names BAD	Alias	BAD; BBC6; BCL2L8; Bcl- XL/Bcl-2 associated death promoter;
Target Names BAD	Product Type	Polyclonal Antibody
	Species	Homo sapiens (Human)
Image	Target Names	BAD
	Image	




CUSABIO TECHNOLOGY LLC










Western blot analysis of extracts from COS7 cells treated with TNF-a using BAD (Phospho-Ser91/128) Antibody. The lane on the right is treated with the antigen-specific peptide.



Immunohistochemical analysis of paraffinembedded human breast carcinoma tissue using BAD (Phospho-Ser91/128) antibody (left)or the same antibody preincubated with blocking peptide (right).

Product Modify

Phospho-Ser91/128