






## SLC22A6 Antibody

Product Code	CSB-PA163258
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q4U2R8
Immunogen	Synthetic peptide of Human SLC22A6
Raised In	Rabbit
Species Reactivity	Human,Rat
<b>Tested Applications</b>	ELISA,WB,IHC;ELISA:1:2000-1:10000,WB:1:1000-1:5000,IHC:1:50-1:200
Relevance	The protein encoded by this gene is involved in the sodium-dependent transport and excretion of organic anions, some of which are potentially toxic. The encoded protein is an integral membrane protein and may be localized to the basolateral membrane. Four transcript variants encoding four different isoforms have been found for this gene. Functions as organic anion exchanger when the uptake of one molecule of organic anion is coupled with an efflux of one molecule of endogenous dicarboxylic acid (glutarate, ketoglutarate, etc). Mediates the sodium-independent uptake of 2,3-dimercapto-1-propanesulfonic acid (DMPS) By similarity.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Purification Method	Antigen affinity purification
Isotype	IgG
Species	Homo sapiens (Human)
Target Names	SLC22A6
Image	The image on the left is immunohistochemistry of




The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using CSB-PA163258(SLC22A6 Antibody) at dilution 1/50, on the right is treated with synthetic peptide. (Original magnification: x200)



## **CUSABIO TECHNOLOGY LLC**







Gel: 10%SDS-PAGE, Lysate: 40 μg, Lane: Human fetal kidney tissue, Primary antibody: CSB-PA163258(SLC22A6 Antibody) at dilution 1/900, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 3 minutes