






## **BCL6** Antibody

<b>Product Code</b>	CSB-PA171912
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P41182
Immunogen	Synthetic peptide of Human BCL6
Raised In	Rabbit
Species Reactivity	Human, Mouse
<b>Tested Applications</b>	ELISA,WB,IHC;ELISA:1:1000-1:2000,WB:1:200-1:1000,IHC:1:25-1:100
Relevance	The protein encoded by this gene is a zinc finger transcription factor and contains an N-terminal POZ domain. This protein acts as a sequence-specific repressor of transcription, and has been shown to modulate the transcription of START-dependent IL-4 responses of B cells. This protein can interact with a variety of POZ-containing proteins that function as transcription corepressors. This gene is found to be frequently translocated and hypermutated in diffuse large-cell lymphoma (DLCL), and may be involved in the pathogenesis of DLCL. Alternatively spliced transcript variants encoding different protein isoforms have been found for this gene
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
<b>Purification Method</b>	Antigen affinity purification
Isotype	IgG
Species	Homo sapiens (Human)
Target Names	BCL6
Image	



The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using CSB-PA171912(BCL6 Antibody) at dilution 1/15, on the right is treated with synthetic peptide. (Original magnification: x200)















The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using CSB-PA171912(BCL6 Antibody) at dilution 1/15, on the right is treated with synthetic peptide. (Original magnification: ×200)



Gel: 6%SDS-PAGE, Lysate: 40 μg, Lane: Raji cells, Primary antibody: CSB-PA171912(BCL6 Antibody) at dilution 1/100, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 10 seconds