


**Image** 





## MAP2K4 (Ab-80) Antibody

<b>Product Code</b>	CSB-PA217719
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P45985
Immunogen	Peptide sequence around aa. 78~82 (T-H-S-I-E) derived from Human SEK1/MKK4.
Raised In	Rabbit
Species Reactivity	Human,Mouse,Rat
Specificity	The antibody detects endogenous level of total SEK1/MKK4 protein.
<b>Tested Applications</b>	ELISA,WB,IHC;WB:1:500-1:1000,IHC:1:50-1:200
Relevance	Dual specificity kinase that activates the JUN kinases MAPK8 (JNK1) and MAPK9 (JNK2) as well as MAPK14 (p38) but not MAPK1 (ERK2) or MAPK3 (ERK1).
	Park HS, et al. (2002) J Biol Chem; 277(4): 2573-8.
Form	Park HS, et al. (2002) J Biol Chem; 277(4): 2573-8.  Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Form Purification Method	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+),
	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-
Purification Method	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitopespecific peptide.
Purification Method  Clonality	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitopespecific peptide.  Polyclonal
Purification Method  Clonality  Alias	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.  Polyclonal  JNKK; JNKK1; MAP2K4; MAPK/ERK kinase 4; kinase 4
Purification Method  Clonality  Alias  Product Type	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.  Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.  Polyclonal  JNKK; JNKK1; MAP2K4; MAPK/ERK kinase 4; kinase 4  Polyclonal Antibody



Western blot analysis of extracts from PC12 and C2C12 cells using SEK1/MKK4(Ab-80) Antibody.




## **CUSABIO TECHNOLOGY LLC**









Immunohistochemical analysis of paraffinembedded human breast carcinoma tissue using SEK1/MKK4(Ab-80) Antibody(left) or the same antibody preincubated with blocking peptide(right).