





## Phospho-PGR (Ser400) Antibody

Product Code	CSB-PA220125		
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.		
Uniprot No.	P06401		
Immunogen	Peptide sequence around phosphorylation site of serine 400 (A-R-S(p)-P-R) derived from Human Progesterone Receptor.		
Raised In	Rabbit		
Species Reactivity	Human		
Specificity	The antibody detects endogenous level of Progesterone Receptor only when phosphorylated at serine 400.		
<b>Tested Applications</b>	ELISA,WB;WB:1:500-1:1000		
Relevance	Progesterone receptors (PRs) are nuclear hormone receptors of the NR3C class, which also includes mineralocorticoid, glucocorticoid and androgen receptors. They exist as homodimers coupled to Hsp90 or HMGB proteins, which are shed upon activation. The major signaling pathway used by progesterone receptors is via direct DNA binding and transcriptional regulation of target genes. They can also signal by binding to other proteins, mainly with transcription factors such as NF-kappaB, AP-1 or STAT. Progesterone receptors are found in the female reproductive tract, mammary glands, brain and pituitary gland and receptor expression is induced by estrogen. Well established functions of progesterone receptors include ovulation, implantation, mammary gland development and maintenance of pregnancy. In addition,progesterone, signaling through the progesterone receptor, increases the ventilatory response of the respiratory centers to carbon dioxide and decreases arterial and alveolar PCO2 in the luteal phase of the menstrual cycle and during pregnancy. The human gene encoding the progesterone receptor has been localized to 11q22.  Hagan CR, Knutson TP, Lange CA (2013) Nucleic Acids Res 41, 8926-42 Wang S, et al. (2013) J Biol Chem 288, 26265-74  Wardell SE, Narayanan R, Weigel NL, Edwards DP (2010) Mol Endocrinol 24, 335-45		
Form	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.		
Purification Method	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi		
Clonality	Polyclonal		
Alias	NR3C3; PGR; PRGR; Progesterone receptor;		



## **CUSABIO TECHNOLOGY LLC**







Product Type	Polyclonal Antibody		
Species	Homo sapiens (Human)		
Target Names	PGR		
Image	KD 170 — 293 Heatshock 130 — 100 — Progesterone Receptor (pSer400) 70 — peptide - +	Western blot analysis of extracts from 293 cells treated with Heatshock using Phospho-Progesterone Receptor (Ser400) antibody. The lane on the right is treated with the antigenspecific peptide.	
<b>Product Modify</b>	Phospho-Ser400		