


Image





CIDEC Antibody

Product Code	CSB-PA244699
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q96AQ7
Immunogen	Fusion protein of Human CIDEC
Raised In	Rabbit
Species Reactivity	Human, Mouse, Rat
Tested Applications	ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:500-1:2000,IHC:1:50-1:200
Relevance	This gene encodes a member of the cell death-inducing DNA fragmentation factor-like effector family. Members of this family play important roles in apoptosis. The encoded protein promotes lipid droplet formation in adipocytes and may mediate adipocyte apoptosis. This gene is regulated by insulin and its expression is positively correlated with insulin sensitivity. Mutations in this gene may contribute to insulin resistant diabetes. A pseudogene of this gene is located on the short arm of chromosome 3. Alternatively spliced transcript variants that encode different isoforms have been observed for this gene.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Purification Method	Antigen affinity purification
Isotype	IgG
Species	Homo sapiens (Human)
Target Names	CIDEC



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using CSB-PA244699(CIDEC Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: ×200)















The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using CSB-PA244699(CIDEC Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: ×200)



Gel: 8%SDS-PAGE, Lysate: 40 μg, Lane 1-2: Mouse intestinum tenue tissue, Mouse heart tissue, Primary antibody: CSB-PA244699(CIDEC Antibody) at dilution 1/551, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 minute