







Phospho-PRKAA1/PRKAA2 (Thr183/Thr172) **Antibody**

Product Code	CSB-PA247163
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Immunogen	Peptide sequence around phosphorylation site of threonine 183(L-R-T(p)-S-C)/threonine172(L- R-T(p)-S-C) derived from Human AMPK?1/AMPK?2.
Raised In	Rabbit
Species Reactivity	Human,Mouse,Rat
Specificity	The antibody detects endogenous level of AMPK α 1/AMPK α 2 only when phosphorylated at threonine 183 or 172.
Tested Applications	ELISA,WB,IHC;WB:1:500-1:1000,IHC:1:50-1:100
Relevance	Catalytic subunit of AMP-activated protein kinase (AMPK), an energy sensor

protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energyproducing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Regulates lipid synthesis by phosphorylating and inactivating lipid metabolic enzymes such as ACACA, ACACB, GYS1, HMGCR and LIPE; regulates fatty acid and cholesterol synthesis by phosphorylating acetyl-CoA carboxylase (ACACA and ACACB) and hormone-sensitive lipase (LIPE) enzymes, respectively. Regulates insulin-signaling and glycolysis by phosphorylating IRS1, PFKFB2 and PFKFB3. AMPK stimulates glucose uptake in muscle by increasing the translocation of the glucose transporter SLC2A4/GLUT4 to the plasma membrane, possibly by mediating phosphorylation of TBC1D4/AS160. Regulates transcription and chromatin structure by phosphorylating transcription regulators involved in energy metabolism such as CRTC2/TORC2, FOXO3, histone H2B, HDAC5, MEF2C, MLXIPL/ChREBP, EP300, HNF4A, p53/TP53, SREBF1, SREBF2 and PPARGC1A. Acts as a key regulator of glucose homeostasis in liver by phosphorylating CRTC2/TORC2, leading to CRTC2/TORC2 sequestration in the cytoplasm. In response to stress, phosphorylates 'Ser-36' of histone H2B (H2BS36ph), leading to promote transcription. Acts as a key regulator of cell growth and proliferation by phosphorylating TSC2, RPTOR and ATG1: in response to nutrient limitation, negatively regulates the mTORC1 complex by phosphorylating RPTOR component of the mTORC1 complex and by phosphorylating and activating TSC2. In response to nutrient limitation, promotes autophagy by phosphorylating and activating ULK1. AMPK also acts as a regulator of circadian rhythm by mediating phosphorylation of CRY1, leading to destabilize it. May regulate the Wnt signaling pathway by phosphorylating CTNNB1, leading to stabilize it. Also has tau-protein kinase









activity: in response to amyloid beta A4 protein (APP) exposure, activated by
CAMKK2, leading to phosphorylation of MAPT/TAU; however the relevance of
such data remains unclear in vivo. Also phosphorylates CFTR, EEF2K, KLC1,
NOS3 and SI C12A1.

Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+),

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide

> and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were

removed by chromatogramphy usi

Clonality Polyclonal

Alias AMPK; AMPKa1; AMPK2


Product Type Polyclonal Antibody

Target Names PRKAA1/PRKAA2


Swiss-Prot: Q13131/P54646; NCBI Gene ID: 5562/5563; NCBI mRNA: Accession NO.

NM_006251.5/ NM_006252.3; NCBI Protein: NP_006242.5 /NP_006243.2

Image



Immunohistochemical analysis of paraffinembedded human lung carcinoma tissue, using AMPK-alpha;1/AMPK-alpha;2(Phospho-Thr174/Thr172) Antibody.



Western blot analysis of extracts from C2C12 cells, treated with serum or calf intestinal phosphatase (CIP), using AMPK α 1/AMPK α 2(Phospho-Thr174/Thr172) Antibody.

Product Modify

Phospho-Thr183/Thr172