


**Image** 





## **FANCG** Antibody

Storage         Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.           Uniprot No.         O15287           Immunogen         Fusion protein of Human FANCG           Raised In         Rabbit           Species Reactivity         Human           Tested Applications         ELISA,IHC;ELISA:1:1000-1:2000,IHC:1:25-1:100           Relevance         The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCJ (also called BRCA2), FANCD4, FANCD4, FANCB, FANCA, FANCB,	<b>Product Code</b>	CSB-PA272939
Immunogen         Fusion protein of Human FANCG           Raised In         Rabbit           Species Reactivity         Human           Tested Applications         ELISA,IHC;ELISA:1:1000-1:2000,IHC:1:25-1:100           Relevance         The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCE, FANCG, FANCI, FANCU (also called BRIP1), FANCL, FANCH and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group G.           Form         Liquid           Conjugate         Non-conjugated           Storage Buffer         -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol           Purification Method         Antigen affinity purification           Isotype         IgG           Species         Homo sapiens (Human)	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISA,IHC;ELISA:1:1000-1:2000,IHC:1:25-1:100RelevanceThe Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCE, FANCE, FANCE, FANCG, FANCI, FANCJ (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group G.FormLiquidConjugateNon-conjugatedStorage Buffer-20°C, pH7.4 PBS, 0.05% NaN3, 40% GlycerolPurification MethodAntigen affinity purificationIsotypeIgGSpeciesHomo sapiens (Human)	Uniprot No.	O15287
Species ReactivityHumanTested ApplicationsELISA,IHC;ELISA:1:1000-1:2000,IHC:1:25-1:100RelevanceThe Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCJ (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group G.FormLiquidConjugateNon-conjugatedStorage Buffer-20°C, pH7.4 PBS, 0.05% NaN3, 40% GlycerolPurification MethodAntigen affinity purificationIsotypeIgGSpeciesHomo sapiens (Human)	Immunogen	Fusion protein of Human FANCG
Tested Applications  ELISA,IHC;ELISA:1:1000-1:2000,IHC:1:25-1:100  Relevance  The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCJ (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group G.  Form  Liquid  Conjugate  Non-conjugated  Storage Buffer  -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol  Purification Method  Antigen affinity purification  Isotype  IgG  Homo sapiens (Human)	Raised In	Rabbit
The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCB, FANCG, FANCI, FANCJ (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group G.  Form Liquid  Conjugate Non-conjugated  Storage Buffer -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol  Purification Method Antigen affinity purification  Isotype IgG  Species Homo sapiens (Human)	Species Reactivity	Human
FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCJ (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group G.  Form Liquid  Conjugate Non-conjugated  Storage Buffer -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol  Purification Method Antigen affinity purification  Isotype IgG  Species Homo sapiens (Human)	<b>Tested Applications</b>	ELISA,IHC;ELISA:1:1000-1:2000,IHC:1:25-1:100
ConjugateNon-conjugatedStorage Buffer-20°C, pH7.4 PBS, 0.05% NaN3, 40% GlycerolPurification MethodAntigen affinity purificationIsotypeIgGSpeciesHomo sapiens (Human)	Relevance	FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCJ (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein
Storage Buffer -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol  Purification Method Antigen affinity purification  Isotype IgG  Species Homo sapiens (Human)	Form	Liquid
Purification Method Antigen affinity purification  Isotype IgG  Species Homo sapiens (Human)	Conjugate	Non-conjugated
Isotype IgG  Species Homo sapiens (Human)	Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Species Homo sapiens (Human)	<b>Purification Method</b>	Antigen affinity purification
· · · · · · · · · · · · · · · · · · ·	Isotype	IgG
Target Names FANCG	Species	Homo sapiens (Human)
	Target Names	FANCG



The image on the left is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using CSB-PA272939(FANCG Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: ×200)




## **CUSABIO TECHNOLOGY LLC**











The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using CSB-PA272939(FANCG Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: ×200)