




LTK Antibody

| Product Code | CSB-PA285877 |
|---|--|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P29376 |
| Immunogen | Synthesized peptide derived from internal of Human LTK. |
| Raised In | Rabbit |
| Species Reactivity | Human, Mouse |
| Specificity | The antibody detects endogenous levels of total LTK protein. |
| Tested Applications | ELISA,WB;WB:1:500-1:3000 |
| Relevance | Orphan receptor with a tyrosine-protein kinase activity. The exact function of this protein is not known. Studies with chimeric proteins (replacing its extracellular region with that of several known growth factor receptors, such as EGFR and CSFIR) demonstrate its ability to promote growth and specifically neurite outgrowth, and cell survival. Signaling appears to involve the PI3 kinase pathway. Involved in regulation of the secretory pathway involving endoplasmic |
| | reticulum (ER) export sites (ERESs) and ER to Golgi transport. Toyoshima H., Proc. Natl. Acad. Sci. U.S.A. 90:5404-5408(1993). Krolewski J.J., EMBO J. 10:2911-2919(1991). Maru Y., Oncogene Res. 5:199-204(1990). |
| Form | Toyoshima H., Proc. Natl. Acad. Sci. U.S.A. 90:5404-5408(1993). Krolewski J.J., EMBO J. 10:2911-2919(1991). |
| Form Purification Method | Toyoshima H., Proc. Natl. Acad. Sci. U.S.A. 90:5404-5408(1993). Krolewski J.J., EMBO J. 10:2911-2919(1991). Maru Y., Oncogene Res. 5:199-204(1990). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, |
| | Toyoshima H., Proc. Natl. Acad. Sci. U.S.A. 90:5404-5408(1993). Krolewski J.J., EMBO J. 10:2911-2919(1991). Maru Y., Oncogene Res. 5:199-204(1990). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity- |
| Purification Method | Toyoshima H., Proc. Natl. Acad. Sci. U.S.A. 90:5404-5408(1993). Krolewski J.J., EMBO J. 10:2911-2919(1991). Maru Y., Oncogene Res. 5:199-204(1990). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Purification Method Clonality | Toyoshima H., Proc. Natl. Acad. Sci. U.S.A. 90:5404-5408(1993). Krolewski J.J., EMBO J. 10:2911-2919(1991). Maru Y., Oncogene Res. 5:199-204(1990). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal EC 2.7.1.112; EC 2.7.10.1; Leukocyte tyrosine kinase receptor precursor; |
| Purification Method Clonality Alias | Toyoshima H., Proc. Natl. Acad. Sci. U.S.A. 90:5404-5408(1993). Krolewski J.J., EMBO J. 10:2911-2919(1991). Maru Y., Oncogene Res. 5:199-204(1990). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal EC 2.7.1.112; EC 2.7.10.1; Leukocyte tyrosine kinase receptor precursor; Protein tyrosine kinase-1; TYK1 |
| Purification Method Clonality Alias Product Type | Toyoshima H., Proc. Natl. Acad. Sci. U.S.A. 90:5404-5408(1993). Krolewski J.J., EMBO J. 10:2911-2919(1991). Maru Y., Oncogene Res. 5:199-204(1990). Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal EC 2.7.1.112; EC 2.7.10.1; Leukocyte tyrosine kinase receptor precursor; Protein tyrosine kinase-1; TYK1 Polyclonal Antibody |



CUSABIO TECHNOLOGY LLC





Western blot analysis of extracts from Jurkat cells, using LTK antibody.