






Phospho-CAMK2A (Thr286) Antibody

Product Code CSB-PA293488 Storage Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Uniprot No. Q9UQM7 Immunogen Peptide sequence around phosphorylation site of threonine 286 (Q-E-T(p)-V-D) derived from Human CaMKII. Raised In Rabbit Species Reactivity Human, Mouse, Rat Specificity The antibody detects endogenous level of CaMKII only when phosphorylated at threonine 286. Tested Applications ELISA,WB;WB:1:500-1:1000 Relevance CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity Pak JH, et al. Proc Natl Acad Sci U S A. 2000 Oct 10; 97(21): 11232-11237 Hudmon A, et al. J Cell Biol. Author manuscript; available in PMC 2006 May 7 Miller P, et al. PLoS Biol. 2005 Apr; 3(4): e107 Runyan JD, et al. Learn Mem. 2005 Mar; 12(2): 103-110. Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clon		
Uniprot No. Q9UQM7 Immunogen Peptide sequence around phosphorylation site of threonine 286 (Q-E-T(p)-V-D) derived from Human CaMKII. Raised In Rabbit Species Reactivity Human,Mouse,Rat Specificity The antibody detects endogenous level of CaMKII only when phosphorylated at threonine 286. Tested Applications ELISA,WB;WB:1:500-1:1000 Relevance CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity Pak JH, et al. Proc Natl Acad Sci U S A. 2000 Oct 10; 97(21): 11232-11237 Hudmon A, et al. J Cell Biol. Author manuscript; available in PMC 2006 May 7 Miller P, et al. PLoS Biol. 2005 Apr; 3(4): e107 Runyan JD, et al. Learn Mem. 2005 Mar; 12(2): 103-110. Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CAMKA Product Type Polyclonal Antibody	Product Code	CSB-PA293488
Immunogen Peptide sequence around phosphorylation site of threonine 286 (Q-E-T(p)-V-D) derived from Human CaMKII. Raised In Rabbit Species Reactivity Human,Mouse,Rat Specificity The antibody detects endogenous level of CaMKII only when phosphorylated at threonine 286. Tested Applications ELISA,WB;WB:1:500-1:1000 Relevance CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity Pak JH, et al. Proc Natl Acad Sci U S A. 2000 Oct 10; 97(21): 11232-11237 Hudmon A, et al. J Cell Biol. Author manuscript; available in PMC 2006 May 7 Miller P, et al. PLoS Biol. 2005 Apr; 3(4): e107 Runyan JD, et al. Learn Mem. 2005 Mar; 12(2): 103-110. Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CAMKA Product Type Polyclonal Antibody Species Homo sapiens (Human)	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
derived from Human CaMKII. Raised In Rabbit Species Reactivity Human,Mouse,Rat Specificity The antibody detects endogenous level of CaMKII only when phosphorylated at threonine 286. Tested Applications ELISA,WB;WB:1:500-1:1000 Relevance CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity Pak JH, et al. Proc Natl Acad Sci U S A. 2000 Oct 10; 97(21): 11232-11237 Hudmon A, et al. J Cell Biol. Author manuscript; available in PMC 2006 May 7 Miller P, et al. PLoS Biol. 2005 Apr; 3(4): e107 Runyan JD, et al. Learn Mem. 2005 Mar; 12(2): 103-110. Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CAMKA Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names CAMK2A	Uniprot No.	Q9UQM7
Specificity Human,Mouse,Rat Specificity The antibody detects endogenous level of CaMKII only when phosphorylated at threonine 286. Tested Applications ELISA,WB;WB:1:500-1:1000 Relevance CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity Pak JH, et al. Proc Natl Acad Sci U S A. 2000 Oct 10; 97(21): 11232-11237 Hudmon A, et al. J Cell Biol. Author manuscript; available in PMC 2006 May 7 Miller P, et al. PLoS Biol. 2005 Apr; 3(4): e107 Runyan JD, et al. Learn Mem. 2005 Mar; 12(2): 103-110. Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CAMKA Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names CAMK2A	Immunogen	
The antibody detects endogenous level of CaMKII only when phosphorylated at threonine 286. Tested Applications ELISA,WB;WB:1:500-1:1000 Relevance CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity Pak JH, et al. Proc Natl Acad Sci U S A. 2000 Oct 10; 97(21): 11232-11237 Hudmon A, et al. J Cell Biol. Author manuscript; available in PMC 2006 May 7 Miller P, et al. PLoS Biol. 2005 Apr; 3(4): e107 Runyan JD, et al. Learn Mem. 2005 Mar; 12(2): 103-110. Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CAMKA Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names CAMK2A	Raised In	Rabbit
Tested Applications ELISA,WB;WB:1:500-1:1000 Relevance CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity Pak JH, et al. Proc Natl Acad Sci U S A. 2000 Oct 10; 97(21): 11232-11237 Hudmon A, et al. J Cell Biol. Author manuscript; available in PMC 2006 May 7 Miller P, et al. PLoS Biol. 2005 Apr; 3(4): e107 Runyan JD, et al. Learn Mem. 2005 Mar; 12(2): 103-110. Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CAMKA Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names CAMK2A	Species Reactivity	Human,Mouse,Rat
Relevance CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity Pak JH, et al. Proc Natl Acad Sci U S A. 2000 Oct 10; 97(21): 11232-11237 Hudmon A, et al. J Cell Biol. Author manuscript; available in PMC 2006 May 7 Miller P, et al. PLoS Biol. 2005 Apr; 3(4): e107 Runyan JD, et al. Learn Mem. 2005 Mar; 12(2): 103-110. Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CAMKA Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names CAMK2A	Specificity	
CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity Pak JH, et al. Proc Natl Acad Sci U S A. 2000 Oct 10; 97(21): 11232-11237 Hudmon A, et al. J Cell Biol. Author manuscript; available in PMC 2006 May 7 Miller P, et al. PLoS Biol. 2005 Apr; 3(4): e107 Runyan JD, et al. Learn Mem. 2005 Mar; 12(2): 103-110. Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CAMKA Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names CAMK2A	Tested Applications	ELISA,WB;WB:1:500-1:1000
Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CAMKA Product Type Polyclonal Antibody Species Homo sapiens (Human) CAMK2A	Relevance	may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity Pak JH, et al. Proc Natl Acad Sci U S A. 2000 Oct 10; 97(21): 11232-11237 Hudmon A, et al. J Cell Biol. Author manuscript; available in PMC 2006 May 7 Miller P, et al. PLoS Biol. 2005 Apr; 3(4): e107
and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CAMKA Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names CAMK2A	Form	
Alias CAMKA Product Type Polyclonal Antibody Species Homo sapiens (Human) Target Names CAMK2A	Purification Method	and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were
Product TypePolyclonal AntibodySpeciesHomo sapiens (Human)Target NamesCAMK2A	Clonality	Polyclonal
Species Homo sapiens (Human) Target Names CAMK2A	Alias	CAMKA
Target Names CAMK2A	Product Type	Polyclonal Antibody
	Species	Homo sapiens (Human)
Image	Target Names	CAMK2A
	Image	




CUSABIO TECHNOLOGY LLC










Western blot analysis of extracts from 293 cells untreated or treated with PMA using CaMKII(Phospho-Thr286) Antibody.



Western blot analysis of extracts from Rat brain tissue treated with Lambda Phosphotase or calf intestinal phosphatase (CIP),using CaMKII (Phospho-Thr286) Antibody.

Product Modify

Phospho-Thr286