


**Image** 





## **ANXA5** Antibody

Product Code	CSB-PA297927
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P08758
Immunogen	Fusion protein of Human ANXA5
Raised In	Rabbit
Species Reactivity	Human,Mouse,Rat
Tested Applications	ELISA,WB,IHC;ELISA:1:2000-1:10000,WB:1:500-1:2000,IHC:1:50-1:200
Relevance	The protein encoded by this gene belongs to the annexin family of calcium-dependent phospholipid binding proteins some of which have been implicated in membrane-related events along exocytotic and endocytotic pathways. Annexin 5 is a phospholipase A2 and protein kinase C inhibitory protein with calcium channel activity and a potential role in cellular signal transduction, inflammation, growth and differentiation. Annexin 5 has also been described as placental anticoagulant protein I, vascular anticoagulant-alpha, endonexin II, lipocortin V, placental protein 4 and anchorin CII. The gene spans 29 kb containing 13 exons, and encodes a single transcript of approximately 1.6 kb and a protein product with a molecular weight of about 35 kDa.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
<b>Purification Method</b>	Antigen affinity purification
Isotype	IgG
Alias	annexin A5
Species	Homo sapiens (Human)
Target Names	ANXA5



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using CSB-PA297927(ANXA5 Antibody) at dilution 1/40, on the right is treated with fusion protein. (Original magnification: ×200)















The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using CSB-PA297927(ANXA5 Antibody) at dilution 1/40, on the right is treated with fusion protein. (Original magnification: ×200)



Gel: 12%SDS-PAGE, Lysate: 40 μg, Lane 1-5: Mouse eyes tissue, Mouse heart tissue, Human cervical cancer tissue, OP9 cells, RAW264.7 cells, Primary antibody: CSB-PA297927(ANXA5 Antibody) at dilution 1/750, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 minute