


Image






TDP2 Antibody

CSB-PA386522
Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
O95551
Synthetic peptide of Human TDP2
Rabbit
Human, Mouse
ELISA,WB,IHC;ELISA:1:1000-1:2000,WB:1:200-1:1000,IHC:1:50-1:200
This gene encodes a member of a superfamily of divalent cation-dependent phosphodiesterases. The encoded protein associates with CD40, tumor necrosis factor (TNF) receptor-75 and TNF receptor associated factors (TRAFs), and inhibits nuclear factor-kappa-B activation. This protein has sequence and structural similarities with APE1 endonuclease, which is involved in both DNA repair and the activation of transcription factors.
Liquid
Non-conjugated
-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Antigen affinity purification
IgG
Homo sapiens (Human)
TDP2



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using CSB-PA386522(TDP2 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: x200)



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using CSB-PA386522(TDP2 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)




CUSABIO TECHNOLOGY LLC









Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane 1-3: Hela cells, 293T cells, mouse liver tissue, Primary antibody: CSB-PA386522(TDP2 Antibody) at dilution 1/200, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure

time: 6 minutes