





MYC Antibody

Product Code	CSB-PA407491
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P01106
Immunogen	Synthetic peptide: C-EQKLISEEDL conjugated with KLH.
Raised In	Rabbit
Species Reactivity	Human,Rat
Tested Applications	ELISA,WB,IHC;ELISA:1:1000-1:10000,WB:1:1000-1:5000,IHC:1:15-1:50
Relevance	A c-myc tag is a polypeptide protein tag derived from the c-myc gene product that can be added to a protein using recombinant DNA technology. It can be used for affinity chromatography, then used to separate recombinant, overexpressed protein from wild type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits.A c-myc tag can be used in many different assays that require recognition by an antibody. If there is no antibody against the studied protein, adding a myc-tag allows one to follow the protein with an antibody against the Myc epitope. Examples are cellular localization studies by immunofluorescence or detection by Western blotting. It can be fused to the C-terminus and the N-terminus of a protein. It is advisable not to fuse the tag directly behind the signal peptide of a secretory protein, since it can interfere with translocation into the secretory pathway.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Purification Method	Antigen affinity purification
Isotype	IgG
Species	Homo sapiens (Human)
Target Names	MYC


Image



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using CSB-PA407491(c-Myc Tag Antibody) at dilution 1/10, on the right is treated with synthetic peptide. (Original magnification: ×200)



The image on the left is immunohistochemistry of paraffin-embedded Human gastric cancer tissue using CSB-PA407491(c-Myc Tag Antibody) at dilution 1/10, on the right is treated with synthetic peptide. (Original magnification: $\times 200$)



Gel: 10%SDS-PAGE, Lysate: 2 μ g, Lane: Recombinant HA tagged protein, Primary antibody: CSB-PA407491(c-Myc Tag Antibody) at dilution 1/1000, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 second