


Image





HIF1AN Antibody

Product Code	CSB-PA447962
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9NWT6
Immunogen	Synthetic peptide of Human HIF1AN
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA,WB,IHC;ELISA:1:1000-1:2000,WB:1:500-1:2000,IHC:1:25-1:100
Relevance	Hypoxia-inducible factor 1-alpha inhibitor is a protein that in humans is encoded by the HIF1AN gene. Hydroxylates HIF-1 alpha at 'Asp-803' in the C-terminal transactivation domain (CAD). Functions as an oxygen sensor and, under normoxic conditions, the hydroxylation prevents interaction of HIF-1 with transcriptional coactivators including Cbp/p300-interacting transactivator. Involved in transcriptional repression through interaction with HIF1A, VHL and histone deacetylases. Hydroxylates specific Asn residues within ankyrin repeat domains (ARD) of NFKB1, NFKBIA, NOTCH1, ASB4, PPP1R12A and several other ARD-containing proteins
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Purification Method	Antigen affinity purification
Isotype	IgG
Species	Homo sapiens (Human)
Target Names	HIF1AN



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using CSB-PA447962(HIF1AN Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: x200)















The image on the left is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using CSB-PA447962(HIF1AN Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: ×200)



Gel: 8%SDS-PAGE, Lysate: 40 µg, Lane: 231 cells, Primary antibody: CSB-PA447962(HIF1AN Antibody) at dilution 1/250, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 20 seconds