


Image





SNAP25 Antibody

| Product Code | CSB-PA564264 |
|----------------------------|--|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P60880 |
| Immunogen | Fusion protein of Human SNAP25 |
| Raised In | Rabbit |
| Species Reactivity | Human,Mouse,Rat |
| Tested Applications | ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:500-1:2000,IHC:1:25-1:100 |
| Relevance | Synaptic vesicle membrane docking and fusion is mediated by SNAREs (soluble N-ethylmaleimide-sensitive factor attachment protein receptors) located on the vesicle membrane (v-SNAREs) and the target membrane (t-SNAREs). The assembled v-SNARE/t-SNARE complex consists of a bundle of four helices, one of which is supplied by v-SNARE and the other three by t-SNARE. For t-SNAREs on the plasma membrane, the protein syntaxin supplies one helix and the protein encoded by this gene contributes the other two. Therefore, this gene product is a presynaptic plasma membrane protein involved in the regulation of neurotransmitter release. Two alternative transcript variants encoding different protein isoforms have been described for this gene. |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol |
| Purification Method | Antigen affinity purification |
| Isotype | IgG |
| Species | Homo sapiens (Human) |
| Target Names | SNAP25 |



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using CSB-PA564264(SNAP25 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: x200)















The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using CSB-PA564264(SNAP25 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)



Gel: 12%SDS-PAGE, Lysate: 40 µg, Lane: Hela cells, Primary antibody: CSB-PA564264(SNAP25 Antibody) at dilution 1/400, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 10 seconds