


Image






CDK20 Antibody

Product Code	CSB-PA578796
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q8IZL9
Immunogen	Fusion protein of Human CDK20
Raised In	Rabbit
Species Reactivity	Human,Mouse,Rat
Tested Applications	ELISA,WB,IHC;ELISA:1:1000-1:2000,WB:1:200-1:1000,IHC:1:25-1:100
Relevance	The protein encoded by this gene contains a kinase domain most closely related to the cyclin-dependent protein kinases. The encoded kinase may activate cyclin-dependent kinase 2 and is involved in cell growth. Alternatively spliced transcript variants encoding distinct isoforms have been reported.
Form	Liquid
Conjugate	Non-conjugated
Conjugate Storage Buffer	Non-conjugated -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
	, ,
Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Storage Buffer Purification Method	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol Antigen affinity purification
Storage Buffer Purification Method Isotype	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol Antigen affinity purification IgG
Storage Buffer Purification Method Isotype Alias	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol Antigen affinity purification IgG cyclin-dependent kinase 20



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using CSB-PA578796(CDK20 Antibody) at dilution 1/15, on the right is treated with fusion protein. (Original magnification: ×200)




The image on the left is immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using CSB-PA578796(CDK20 Antibody) at dilution 1/15, on the right is treated with fusion protein. (Original magnification: x200)



CUSABIO TECHNOLOGY LLC







Gel: 8%SDS-PAGE, Lysate: 40 μ g, Lane: HT29 cells, Primary antibody: CSB-PA578796(CDK20 Antibody) at dilution 1/300, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 5 minutes