


Image






PIK3CD Antibody

CSB-PA597940
Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
O00329
Synthetic peptide of Human PIK3CD
Rabbit
Human, Mouse
ELISA,IHC;ELISA:1:2000-1:5000,IHC:1:50-1:200
Phosphoinositide 3-kinases (PI3Ks) phosphorylate inositol lipids and are involved in the immune response. The protein encoded by this gene is a class I PI3K found primarily in leukocytes. Like other class I PI3Ks (p110-alpha p110-beta, and p110-gamma), the encoded protein binds p85 adapter proteins and GTP-bound RAS. However, unlike the other class I PI3Ks, this protein phosphorylates itself, not p85 protein.
Liquid
Non-conjugated
-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Antigen affinity purification
IgG
Homo sapiens (Human)
PIK3CD



The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using CSB-PA597940(PIK3CD Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: ×200)



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using CSB-PA597940(PIK3CD Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: ×200)