

**Image** 









## **ARMCX2** Antibody

| CSB-PA617569   |
|--|
| Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| Q7L311   |
| Fusion protein of Human ARMCX2   |
| Rabbit   |
| Human, Mouse   |
| ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:500-1:2000,IHC:1:50-1:200  |
| This gene encodes a protein containing a potential N-terminal transmembrane domain and multiple armadillo (arm) repeats. Proteins containing arm repeats are involved in development, maintenance of tissue integrity, and tumorigenesis. This gene is located in a cluster of related genes on chromosome X. There is a pseudogene for this gene on chromosome 7. Alternative splicing in the 5' UTR results in multiple transcript variants encoding the same protein. |
| Liquid   |
| Non-conjugated   |
| -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol   |
| Antigen affinity purification  |
| IgG  |
| Homo sapiens (Human)   |
| ARMCX2   |
|  |



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using CSB-PA617569(ARMCX2 Antibody) at dilution 1/40, on the right is treated with fusion protein. (Original magnification: ×200)




The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using CSB-PA617569(ARMCX2 Antibody) at dilution 1/40, on the right is treated with fusion protein. (Original magnification: ×200)



## **CUSABIO TECHNOLOGY LLC**







Gel: 6%SDS-PAGE, Lysate: 40  $\mu g$ , Lane: Mouse testis tissue, Primary antibody: CSB-PA617569(ARMCX2 Antibody) at dilution 1/425, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 40 seconds