





NEDD8 Antibody

Product Code	CSB-PA618018ESR1HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q15843
Immunogen	Recombinant Human NEDD8 protein (1-81AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200, IF:1:50-1:200
Relevance	Ubiquitin-like protein which plays an important role in cell cycle control and embryogenesis. Covalent attachment to its substrates requires prior activation by the E1 complex UBE1C-APPBP1 and linkage to the E2 enzyme UBE2M. Attachment of NEDD8 to cullins activates their associated E3 ubiquitin ligase activity, and thus promotes polyubiquitination and proteasomal degradation of cyclins and other regulatory proteins.
Form	Liquid
Form Conjugate	
	Liquid
Conjugate	Liquid Non-conjugated
Conjugate Storage Buffer	Liquid Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Conjugate Storage Buffer Purification Method	Liquid Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Antigen Affinity Purified
Conjugate Storage Buffer Purification Method Isotype	Liquid Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Antigen Affinity Purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	Liquid Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Antigen Affinity Purified IgG Polyclonal NEDD8 (Neddylin) (Neural precursor cell expressed developmentally down-
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	Liquid Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Antigen Affinity Purified IgG Polyclonal NEDD8 (Neddylin) (Neural precursor cell expressed developmentally down-regulated protein 8) (NEDD-8) (Ubiquitin-like protein Nedd8), NEDD8
Conjugate Storage Buffer Purification Method Isotype Clonality Alias Species	Liquid Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Antigen Affinity Purified IgG Polyclonal NEDD8 (Neddylin) (Neural precursor cell expressed developmentally down-regulated protein 8) (NEDD-8) (Ubiquitin-like protein Nedd8), NEDD8 Human



20kDa

Western blot

All lanes: NEDD8 antibody at 16µg/ml

Lane 1: Mouse heart tissue Lane 2: Mouse spleen tissue


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 9 kDa Observed band size: 9 kDa










Immunohistochemistry of paraffin-embedded human breast cancer using CSB-PA618018ESR1HU at dilution of 1:100



Immunofluorescent analysis of Hela cells using CSB-PA618018ESR1HU at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)