





MERTK Antibody

| Storage Uniprot No. Immunogen Raised In Species Reactivity | CSB-PA621519ESR1HU Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Q12866 Recombinant Human Tyrosine-protein kinase Mer protein (840-999AA) Rabbit Human, Mouse |
|--|---|
| Uniprot No. Immunogen Raised In Species Reactivity | Q12866 Recombinant Human Tyrosine-protein kinase Mer protein (840-999AA) Rabbit |
| Immunogen Raised In Species Reactivity | Recombinant Human Tyrosine-protein kinase Mer protein (840-999AA) Rabbit |
| Raised In Species Reactivity | Rabbit |
| Species Reactivity | |
| | Human, Mouse |
| Tested Applications | |
| • • • | ELISA, WB; Recommended dilution: WB:1:200-1:1000 |
| | Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to several ligands including LGALS3, TUB, TULP1 or GAS6. Regulates many physiological processes including cell survival, migration, differentiation, and phagocytosis of apoptotic cells (efferocytosis). Ligand binding at the cell surface induces autophosphorylation of MERTK on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with GRB2 or PLCG2 and induces phosphorylation of MAPK1, MAPK2, FAK/PTK2 or RAC1. MERTK signaling plays a role in various processes such as macrophage clearance of apoptotic cells, platelet aggregation, cytoskeleton reorganization and engulfment. Functions in the retinal pigment epithelium (RPE) as a regulator of rod outer segments fragments phagocytosis. Plays also an important role in inhibition of Toll-like receptors (TLRs)-mediated innate immune response by activating STAT1, which selectively induces production of suppressors of cytokine signaling SOCS1 and SOCS3. |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | PBS with 0.02% sodium azide, 50% glycerol, pH7.3. |
| Purification Method | Antigen Affinity Purified |
| Isotype | IgG |
| | |
| | Polyclonal |
| Clonality Alias | Polyclonal Tyrosine-protein kinase Mer (EC 2.7.10.1) (Proto-oncogene c-Mer) (Receptor tyrosine kinase MerTK), MERTK, MER |
| Clonality Alias | Tyrosine-protein kinase Mer (EC 2.7.10.1) (Proto-oncogene c-Mer) (Receptor |
| Clonality Alias Species | Tyrosine-protein kinase Mer (EC 2.7.10.1) (Proto-oncogene c-Mer) (Receptor tyrosine kinase MerTK), MERTK, MER |
| Clonality Alias Species Research Area | Tyrosine-protein kinase Mer (EC 2.7.10.1) (Proto-oncogene c-Mer) (Receptor tyrosine kinase MerTK), MERTK, MER Human |




CUSABIO TECHNOLOGY LLC









Western blot

All lanes: MERTK antibody at 8μg/ml

Lane 1: Mouse gonad tissue Lane 2: Mouse lung tissue Lane 3: Mouse kidney tissue

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 110 kDa Observed band size: 110 kDa