





KCNMB1 Antibody, FITC conjugated

Product Code	CSB-PA621958LC01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q16558
Immunogen	Recombinant Human Calcium-activated potassium channel subunit beta-1 protein (40-130AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	Regulatory subunit of the calcium activated potassium KCNMA1 (maxiK) channel. Modulates the calcium sensitivity and gating kinetics of KCNMA1, thereby contributing to KCNMA1 channel diversity. Increases the apparent Ca(2+)/voltage sensitivity of the KCNMA1 channel. It also modifies KCNMA1 channel kinetics and alters its pharmacological properties. It slows down the activation and the deactivation kinetics of the channel. Acts as a negative regulator of smooth muscle contraction by enhancing the calcium sensitivity to KCNMA1. Its presence is also a requirement for internal binding of the KCNMA1 channel opener dehydrosoyasaponin I (DHS-1) triterpene glycoside and for external binding of the agonist hormone 17-beta-estradiol (E2). Increases the
	binding activity of charybdotoxin (CTX) toxin to KCNMA1 peptide blocker by increasing the CTX association rate and decreasing the dissociation rate.
Form	
Form Conjugate	increasing the CTX association rate and decreasing the dissociation rate.
	increasing the CTX association rate and decreasing the dissociation rate. Liquid
Conjugate	increasing the CTX association rate and decreasing the dissociation rate. Liquid FITC Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	increasing the CTX association rate and decreasing the dissociation rate. Liquid FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	increasing the CTX association rate and decreasing the dissociation rate. Liquid FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Conjugate Storage Buffer Purification Method Isotype	increasing the CTX association rate and decreasing the dissociation rate. Liquid FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	increasing the CTX association rate and decreasing the dissociation rate. Liquid FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Calcium-activated potassium channel subunit beta-1 (BK channel subunit beta-1) (BKbeta) (BKbeta1) (Hbeta1) (Calcium-activated potassium channel, subfamily M subunit beta-1) (Calcium-activated potassium channel subunit beta) (Charybdotoxin receptor subunit beta-1) (K(VCA)beta-1) (Maxi K channel
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	increasing the CTX association rate and decreasing the dissociation rate. Liquid FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Calcium-activated potassium channel subunit beta-1 (BK channel subunit beta-1) (BKbeta) (BKbeta1) (Hbeta1) (Calcium-activated potassium channel, subfamily M subunit beta-1) (Calcium-activated potassium channel subunit beta) (Charybdotoxin receptor subunit beta-1) (K(VCA)beta-1) (Maxi K channel subunit beta-1) (Slo-beta-1) (Slo-beta), KCNMB1