





## TRIP13 Antibody


CSB-PA623008LA01HU
Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Q15645
Recombinant Human Pachytene checkpoint protein 2 homolog protein (181-390AA)
Rabbit
Human
ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200
Plays a key role in chromosome recombination and chromosome structure development during meiosis. Required at early steps in meiotic recombination that leads to non-crossovers pathways. Also needed for efficient completion of homologous synapsis by influencing crossover distribution along the chromosomes affecting both crossovers and non-crossovers pathways. Also required for development of higher-order chromosome structures and is needed for synaptonemal-complex formation. In males, required for efficient synapsis of the sex chromosomes and for sex body formation. Promotes early steps of the DNA double-strand breaks (DSBs) repair process upstream of the assembly of RAD51 complexes. Required for depletion of HORMAD1 and HORMAD2 from synapsed chromosomes (By similarity). Plays a role in mitotic spindle assembly
checkpoint (SAC) activation (PubMed:28553959).
checkpoint (SAC) activation (PubMed:28553959).  Liquid
, , , , , , , , , , , , , , , , , , , ,
Liquid
Liquid Non-conjugated Preservative: 0.03% Proclin 300
Liquid  Non-conjugated  Preservative: 0.03% Proclin 300  Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Liquid  Non-conjugated  Preservative: 0.03% Proclin 300  Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Antigen Affinity Purified
Liquid  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Antigen Affinity Purified  IgG
Liquid  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Antigen Affinity Purified  IgG  Polyclonal  Pachytene checkpoint protein 2 homolog (Human papillomavirus type 16 E1 protein-binding protein) (16E1-BP) (HPV16 E1 protein-binding protein) (Thyroid hormone receptor interactor 13) (Thyroid receptor-interacting protein 13) (TR-
Liquid  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Antigen Affinity Purified  IgG  Polyclonal  Pachytene checkpoint protein 2 homolog (Human papillomavirus type 16 E1 protein-binding protein) (16E1-BP) (HPV16 E1 protein-binding protein) (Thyroid hormone receptor interactor 13) (Thyroid receptor-interacting protein 13) (TR-interacting protein 13) (TRIP-13), TRIP13, PCH2
Liquid  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Antigen Affinity Purified  IgG  Polyclonal  Pachytene checkpoint protein 2 homolog (Human papillomavirus type 16 E1 protein-binding protein) (16E1-BP) (HPV16 E1 protein-binding protein) (Thyroid hormone receptor interactor 13) (Thyroid receptor-interacting protein 13) (TR-interacting protein 13) (TRIP-13), TRIP13, PCH2













Immunohistochemistry of paraffin-embedded human testis tissue using CSB-PA623008LA01HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human placenta tissue using CSB-PA623008LA01HU at dilution of 1:100



Western blot

All lanes: TRIP13 antibody at 1µg/ml + A549

whole cell lysate Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 49, 23 kDa Observed band size: 49 kDa