





HSD17B12 Antibody, FITC conjugated

| Product Code | CSB-PA687494LC01HU |
|----------------------------|--|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q53GQ0 |
| Immunogen | Recombinant Human Very-long-chain 3-oxoacyl-CoA reductase protein (40-181AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA |
| Relevance | Catalyzes the second of the four reactions of the long-chain fatty acids elongation cycle. This endoplasmic reticulum-bound enzymatic process, allows the addition of two carbons to the chain of long- and very long-chain fatty acids/VLCFAs per cycle. This enzyme has a 3-ketoacyl-CoA reductase activity, reducing 3-ketoacyl-CoA to 3-hydroxyacyl-CoA, within each cycle of fatty acid elongation. Thereby, it may participate to the production of VLCFAs of different chain lengths that are involved in multiple biological processes as precursors of membrane lipids and lipid mediators. May also catalyze the transformation of estrone (E1) into estradiol (E2) and play a role in estrogen formation. |
| Form | Liquid |
| Conjugate | FITC |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Purification Method | Antigen Affinity Purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Very-long-chain 3-oxoacyl-CoA reductase (EC 1.1.1.330) (17-beta-hydroxysteroid dehydrogenase 12) (17-beta-HSD 12) (3-ketoacyl-CoA reductase) (KAR) (Estradiol 17-beta-dehydrogenase 12) (EC 1.1.1.62) (Short chain dehydrogenase/reductase family 12C member 1), HSD17B12, SDR12C1 |
| Species | Human |
| Research Area | |
| Research Area | Signal Transduction |