


Image





TGFB1 Antibody

CSB-PA697937
Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
P01137
Fusion protein of Human TGFB1
Rabbit
Human, Mouse, Rat
ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:500-1:2000,IHC:1:25-1:100
This gene encodes a member of the transforming growth factor beta (TGFB) family of cytokines, which are multifunctional peptides that regulate proliferation, differentiation, adhesion, migration, and other functions in many cell types. Many cells have TGFB receptors, and the protein positively and negatively regulates many other growth factors. The secreted protein is cleaved into a latency-associated peptide (LAP) and a mature TGFB1 peptide, and is found in either a latent form composed of a TGFB1 homodimer, a LAP homodimer, and a latent TGFB1-binding protein, or in an active form composed of a TGFB1 homodimer. The mature peptide may also form heterodimers with other TGFB family members. This gene is frequently upregulated in tumor cells, and mutations in this gene result in Camurati-Engelmann disease.
Liquid
Non-conjugated
-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Antigen affinity purification
IgG
Homo sapiens (Human)
TGFB1



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using CSB-PA697937(TGFB1 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: X200)















The image on the left is immunohistochemistry of paraffin-embedded Human lung cancer tissue using CSB-PA697937(TGFB1 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: X200)



Gel: 10%SDS-PAGE, Lysate: 40 µg, Lane: Mouse heart tissue, Primary antibody: CSB-PA697937(TGFB1 Antibody) at dilution 1/450, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 20 seconds