


Image





ATG16L1 Antibody


Product Code	CSB-PA734569LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q676U5
Immunogen	Recombinant Human Autophagy-related protein 16-1 protein (1-607AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:200-1:500
Relevance	Plays an essential role in autophagy: interacts with ATG12-ATG5 to mediate the conjugation of phosphatidylethanolamine (PE) to LC3 (MAP1LC3A, MAP1LC3B or MAP1LC3C), to produce a membrane-bound activated form of LC3 named LC3-II. Thereby, controls the elongation of the nascent autophagosomal membrane.
Form	Liquid
Form Conjugate	Liquid Non-conjugated
Conjugate	Non-conjugated Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Conjugate Storage Buffer Purification Method Isotype	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Autophagy-related protein 16-1 (APG16-like 1), ATG16L1, APG16L
Conjugate Storage Buffer Purification Method Isotype Clonality Alias Species	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Autophagy-related protein 16-1 (APG16-like 1), ATG16L1, APG16L Human



Immunofluorescence staining of HepG2 cells with CSB-PA734569LA01HU at 1:333, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).




CUSABIO TECHNOLOGY LLC












Immunohistochemistry of paraffin-embedded human skin tissue using CSB-PA734569LA01HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human colon cancer using CSB-PA734569LA01HU at dilution of 1:100