

🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com


## **RRM2B** Antibody

| Product Code               | CSB-PA745339ESR2HU   |  |
|----------------------------|--|--|
| Storage                    | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |  |
| Uniprot No.                | Q7LG56   |  |
| Immunogen                  | Recombinant Human Ribonucleoside-diphosphate reductase subunit M2 B protein (1-351AA)  |  |
| Raised In                  | Rabbit   |  |
| Species Reactivity         | Human  |  |
| <b>Tested Applications</b> | ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200  |  |
| Relevance                  | Plays a pivotal role in cell survival by repairing damaged DNA in a p53/TP53-<br>dependent manner. Supplies deoxyribonucleotides for DNA repair in cells<br>arrested at G1 or G2. Contains an iron-tyrosyl free radical center required for<br>catalysis. Forms an active ribonucleotide reductase (RNR) complex with RRM1<br>which is expressed both in resting and proliferating cells in response to DNA<br>damage. |  |
| Form                       | Liquid   |  |
| Conjugate                  | Non-conjugated   |  |
| Storage Buffer             | PBS with 0.02% sodium azide, 50% glycerol, pH7.3.  |  |
| Purification Method        | Antigen Affinity Purified  |  |
| Isotype                    | lgG  |  |
| Clonality                  | Polyclonal   |  |
| Alias                      | Ribonucleoside-diphosphate reductase subunit M2 B (EC 1.17.4.1) (TP53-<br>inducible ribonucleotide reductase M2 B) (p53-inducible ribonucleotide<br>reductase small subunit 2-like protein) (p53R2), RRM2B, P53R2  |  |
| Species                    | Human  |  |
| Research Area              | Epigenetics and Nuclear Signaling  |  |
| Target Names               | RRM2B  |  |
| Image                      | 120kDa —<br>90kDa —<br>50kDa —<br>35kDa —<br>25kDa —<br>20kDa —  | Western blot<br>All lanes: RRM2B antibody at 3µg/ml<br>Lane 1: MCF-7 whole cell lysate<br>Lane 2: Hela whole cell lysate<br>Lane 3: HepG2 whole cell lysate<br>Secondary<br>Goat polyclonal to rabbit IgG at 1/10000 dilution<br>Predicted band size: 41, 35, 16, 8, 5, 49 kDa<br>Observed band size: 41 kDa |

1



🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🥥 Website: www.cusabio.com 🥣



Immunohistochemistry of paraffin-embedded human tonsil tissue using CSB-PA745339ESR2HU at dilution of 1:100