




PLCZ1 Antibody

Product Code	CSB-PA768240LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q86YW0
Immunogen	Recombinant Human 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase zeta-1 protein (1-415AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB; Recommended dilution: WB:1:1000-1:5000
Relevance	The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. In vitro, hydrolyzes PtdIns(4,5)P2 in a Ca(2+)-dependent manner. Triggers intracellular Ca(2+) oscillations in oocytes solely during M phase and is involved in inducing oocyte activation and initiating embryonic development up to the blastocyst stage. Is therefore a strong candidate for the egg-activating soluble sperm factor that is transferred from the sperm into the egg cytoplasm following gamete membrane fusion. May exert an inhibitory effect on phospholipase-C-coupled processes that depend on calcium ions and protein kinase C, including CFTR trafficking and function.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase zeta-1 (EC 3.1.4.11) (Phosphoinositide phospholipase C-zeta-1) (Phospholipase C-zeta-1) (PLC-zeta-1) (Testis-development protein NYD-SP27), PLCZ1
Species	Human
Research Area	Cancer
Target Names	PLCZ1
Image	



Western blot

All lanes: PLCZ1 antibody at 3 μ g/ml

Lane 1: Mouse liver tissue

Lane 2: Mouse kidney tissue

Lane 3: Human placenta tissue

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 71, 48, 58 kDa

Observed band size: 71 kDa