

Image








LIMK1 Antibody

| Product Code | CSB-PA782037 |
|----------------------------|--|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P53667 |
| Immunogen | Synthetic peptide of Human LIMK1 |
| Raised In | Rabbit |
| Species Reactivity | Human,Mouse,Rat |
| Tested Applications | ELISA,IHC;ELISA:1:2000-1:5000,IHC:1:50-1:200 |
| Relevance | There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. LIMK1 is a serine/threonine kinase that regulates actin polymerization via phosphorylation and inactivation of the actin binding factor cofilin. This protein is ubiquitously expressed during development and plays a role in many cellular processes associated with cytoskeletal structure. |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol |
| Purification Method | Antigen affinity purification |
| Isotype | IgG |
| Species | Homo sapiens (Human) |
| Target Names | LIMK1 |



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using CSB-PA782037(LIMK1 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)




CUSABIO TECHNOLOGY LLC











The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using CSB-PA782037(LIMK1 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)