






IL22RA1 Antibody

Product Code	CSB-PA822717LA01HU
Abbreviation	Interleukin-22 receptor subunit alpha-1
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q8N6P7
Immunogen	Recombinant Human Interleukin-22 receptor subunit alpha-1 protein (250-573AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:500
Relevance	Component of the receptor for IL20, IL22 and IL24. Component of IL22 receptor formed by IL22RA1 and IL10RB enabling IL22 signaling via JAK/STAT pathways. IL22 also induces activation of MAPK1/MAPK3 and Akt kinases pathways. Component of one of the receptor for IL20 and IL24 formed by IL22RA1 and IL20RB also signaling through STATs activation. Mediates IL24 antiangiogenic activity as well as IL24 inhibitory effect on endothelial cell tube formation and differentiation.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	
Furnication Method	>95%, Protein G purified
Isotype	>95%, Protein G purified IgG
	· · · · · · · · · · · · · · · · · · ·
Isotype	IgG
Isotype Clonality	IgG Polyclonal Interleukin-22 receptor subunit alpha-1 (IL-22 receptor subunit alpha-1) (IL-22R-alpha-1) (IL-22RA1) (Cytokine receptor class-II member 9) (Cytokine receptor
Isotype Clonality Alias	IgG Polyclonal Interleukin-22 receptor subunit alpha-1 (IL-22 receptor subunit alpha-1) (IL-22R-alpha-1) (IL-22RA1) (Cytokine receptor class-II member 9) (Cytokine receptor family 2 member 9) (CRF2-9) (ZcytoR11), IL22RA1, IL22R
Isotype Clonality Alias Species	Polyclonal Interleukin-22 receptor subunit alpha-1 (IL-22 receptor subunit alpha-1) (IL-22R-alpha-1) (IL-22RA1) (Cytokine receptor class-II member 9) (Cytokine receptor family 2 member 9) (CRF2-9) (ZcytoR11), IL22RA1, IL22R Human



Immunofluorescence staining of HepG2 cells with CSB-PA822717LA01HU at 1:333, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).




CUSABIO TECHNOLOGY LLC









Immunohistochemistry of paraffin-embedded human tonsil tissue using CSB-PA822717LA01HU at dilution of 1:100