





PNPLA2 Antibody, FITC conjugated

Product Code	CSB-PA836180LC01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q96AD5
Immunogen	Recombinant Human Patatin-like phospholipase domain-containing protein 2 protein (1-504AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	Catalyzes the initial step in triglyceride hydrolysis in adipocyte and non-adipocyte lipid droplets (PubMed:15550674). Also has acylglycerol transacylase activity. May act coordinately with LIPE/HLS within the lipolytic cascade. Regulates adiposome size and may be involved in the degradation of adiposomes (PubMed:16239926). May play an important role in energy homeostasis. May play a role in the response of the organism to starvation, enhancing hydrolysis of triglycerides and providing free fatty acids to other tissues to be exidized in cituations of energy depletion.
	tissues to be oxidized in situations of energy depletion.
Form	Liquid
Form Conjugate	• • • • • • • • • • • • • • • • • • • •
	Liquid
Conjugate	Liquid FITC Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	Liquid FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	Liquid FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Conjugate Storage Buffer Purification Method Isotype	Liquid FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	Liquid FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Patatin-like phospholipase domain-containing protein 2 (EC 3.1.1.3) (Adipose triglyceride lipase) (Calcium-independent phospholipase A2) (Desnutrin) (IPLA2-zeta) (Pigment epithelium-derived factor) (TTS2.2) (Transport-secretion protein
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	Liquid FITC Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Patatin-like phospholipase domain-containing protein 2 (EC 3.1.1.3) (Adipose triglyceride lipase) (Calcium-independent phospholipase A2) (Desnutrin) (IPLA2-zeta) (Pigment epithelium-derived factor) (TTS2.2) (Transport-secretion protein 2) (TTS2), PNPLA2, ATGL