




RNF128 Antibody

Product Code	CSB-PA837448LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q8TEB7
Immunogen	Recombinant Human E3 ubiquitin-protein ligase RNF128 protein (229-400AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB, IF; Recommended dilution: WB:1:1000-1:5000, IF:1:50-1:200
Relevance	E3 ubiquitin-protein ligase that catalyzes 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains formation. Functions as an inhibitor of cytokine gene transcription. Inhibits IL2 and IL4 transcription, thereby playing an important role in the induction of the anergic phenotype, a long-term stable state of T-lymphocyte unresponsiveness to antigenic stimulation associated with the blockade of interleukin production. Ubiquitinates ARPC5 with 'Lys-48' linkages and COR1A with 'Lys-63' linkages leading to their degradation, down-regulation of these cytoskeletal components results in impaired lamellipodium formation and reduced accumulation of F-actin at the immunological synapse. Functions in the patterning of the dorsal ectoderm; sensitizes ectoderm to respond to neural-inducing signals.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	E3 ubiquitin-protein ligase RNF128 (EC 2.3.2.27) (Gene related to anergy in lymphocytes protein) (GRAIL) (RING finger protein 128) (RING-type E3 ubiquitin transferase RNF128), RNF128
Species	Human
Research Area	Cell Biology
Target Names	RNF128
Image	



Western blot


All lanes: RNF128 antibody at 12 μ g/ml + Mouse liver tissue

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 47, 45 kDa

Observed band size: 47 kDa



Immunofluorescent analysis of HepG2 cells

using CSB-PA837448LA01HU at dilution of

1:100 and Alexa Fluor 488-conjugated

AffiniPure Goat Anti-Rabbit IgG(H+L)