





HSD17B11 Antibody, HRP conjugated

| Product Code | CSB-PA843318HB01HU |
|----------------------------|---|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q8NBQ5 |
| Immunogen | Recombinant Human Estradiol 17-beta-dehydrogenase 11 protein (20-300AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA |
| Relevance | Can convert androstan-3-alpha,17-beta-diol (3-alpha-diol) to androsterone in vitro, suggesting that it may participate in androgen metabolism during steroidogenesis. May act by metabolizing compounds that stimulate steroid synthesis and/or by generating metabolites that inhibit it. Has no activity toward DHEA (dehydroepiandrosterone), or A-dione (4-androste-3,17-dione), and only a slight activity toward testosterone to A-dione. Tumor-associated antigen in cutaneous T-cell lymphoma. |
| Form | Liquid |
| Conjugate | HRP |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Estradiol 17-beta-dehydrogenase 11 (EC 1.1.1.62) (17-beta-hydroxysteroid dehydrogenase 11) (17-beta-HSD 11) (17bHSD11) (17betaHSD11) (17-beta-hydroxysteroid dehydrogenase XI) (17-beta-HSD XI) (17betaHSDXI) (Cutaneous T-cell lymphoma-associated antigen HD-CL-03) (CTCL-associated antigen HD-CL-03) (Dehydrogenase/reductase SDR family member 8) (Retinal short-chain dehydrogenase/reductase 2) (retSDR2) (Short chain dehydrogenase/reductase family 16C member 2), HSD17B11, DHRS8 PAN1B SDR16C2 |
| Species | Homo sapiens (Human) |
| Research Area | Signal Transduction |
| Target Names | HSD17B11 |