



HSD17B11 Antibody, FITC conjugated

Product Code	CSB-PA843318HC01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q8NBQ5
Immunogen	Recombinant Human Estradiol 17-beta-dehydrogenase 11 protein (20-300AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	Can convert androstan-3-alpha,17-beta-diol (3-alpha-diol) to androsterone in vitro, suggesting that it may participate in androgen metabolism during steroidogenesis. May act by metabolizing compounds that stimulate steroid synthesis and/or by generating metabolites that inhibit it. Has no activity toward DHEA (dehydroepiandrosterone), or A-dione (4-androste-3,17-dione), and only a slight activity toward testosterone to A-dione. Tumor-associated antigen in cutaneous T-cell lymphoma.
Form	Liquid
Conjugate	FITC
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Estradiol 17-beta-dehydrogenase 11 (EC 1.1.1.62) (17-beta-hydroxysteroid dehydrogenase 11) (17-beta-HSD 11) (17bHSD11) (17betaHSD11) (17-beta-hydroxysteroid dehydrogenase XI) (17-beta-HSD XI) (17betaHSDXI) (Cutaneous T-cell lymphoma-associated antigen HD-CL-03) (CTCL-associated antigen HD-CL-03) (Dehydrogenase/reductase SDR family member 8) (Retinal short-chain dehydrogenase/reductase 2) (retSDR2) (Short chain dehydrogenase/reductase family 16C member 2), HSD17B11, DHRS8 PAN1B SDR16C2
Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	HSD17B11