






SMARCA4 Antibody

Product Code	CSB-PA843993
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P51532
Immunogen	Synthetic peptide of Human SMARCA4
Raised In	Rabbit
Species Reactivity	Human,Mouse,Rat
Tested Applications	ELISA,WB,IHC;ELISA:1:1000-1:5000,WB:1:200-1:1000,IHC:1:50-1:200
Relevance	The protein encoded by this gene is a member of the SWI/SNF family of proteins and is similar to the brahma protein of Drosophila. Members of this family have helicase and ATPase activities and are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent chromatin remodeling complex SNF/SWI, which is required for transcriptional activation of genes normally repressed by chromatin. In addition, this protein can bind BRCA1, as well as regulate the expression of the tumorigenic protein CD44. Mutations in this gene cause rhabdoid tumor predisposition syndrome type 2. Multiple transcript variants encoding different isoforms have been found for this gene.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Purification Method	Antigen affinity purification
Isotype	IgG
Species	Homo sapiens (Human)
Target Names	SMARCA4
Image	The image on the left is immunohistochemistry of




The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using CSB-PA843993(SMARCA4 Antibody) at dilution 1/50, on the right is treated with synthetic peptide. (Original magnification: ×200)












The image on the left is immunohistochemistry of paraffin-embedded Human lung cancer tissue using CSB-PA843993(SMARCA4 Antibody) at dilution 1/50, on the right is treated with synthetic peptide. (Original magnification: ×200)



Gel: 8%SDS-PAGE, Lysate: 50 µg, Lane 1-3: K562 cells, 293T cells, hepG2 cells, Primary antibody: CSB-PA843993(SMARCA4 Antibody) at dilution 1/300, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 1 minute