





## SCARB1 Antibody

<b>Product Code</b>	CSB-PA845139LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q8WTV0
Immunogen	Recombinant Human Scavenger receptor class B member 1 protein (33-443AA)
Raised In	Rabbit
Species Reactivity	Human
<b>Tested Applications</b>	ELISA, IHC; Recommended dilution: IHC:1:20-1:200
Relevance	Receptor for different ligands such as phospholipids, cholesterol ester, lipoproteins, phosphatidylserine and apoptotic cells. Probable receptor for HDL, located in particular region of the plasma membrane, called caveolae. Facilitates the flux of free and esterified cholesterol between the cell surface and extracellular donors and acceptors, such as HDL and to a lesser extent, apoB-containing lipoproteins and modified lipoproteins. Probably involved in the phagocytosis of apoptotic cells, via its phosphatidylserine binding activity. Receptor for hepatitis C virus glycoprotein E2. Binding between SCARB1 and
	E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester.
Form	E2 was found to be independent of the genotype of the viral isolate. Plays an
Form Conjugate	E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester .
	E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester .  Liquid
Conjugate	E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester .  Liquid  Non-conjugated  Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester .  Liquid  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester .  Liquid  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified
Conjugate Storage Buffer Purification Method Isotype	E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester .  Liquid  Non-conjugated  Preservative: 0.03% Proclin 300  Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified  IgG
Conjugate Storage Buffer  Purification Method Isotype Clonality	E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester .  Liquid  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified  IgG  Polyclonal  Scavenger receptor class B member 1 (SRB1) (CD36 and LIMPII analogous 1) (CLA-1) (CD36 antigen-like 1) (Collagen type I receptor, thrombospondin
Conjugate Storage Buffer  Purification Method Isotype Clonality Alias	E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester .  Liquid  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified  IgG  Polyclonal  Scavenger receptor class B member 1 (SRB1) (CD36 and LIMPII analogous 1) (CLA-1) (CD36 antigen-like 1) (Collagen type I receptor, thrombospondin receptor-like 1) (SR-BI) (CD antigen CD36), SCARB1, CD36L1 CLA1
Conjugate Storage Buffer  Purification Method Isotype Clonality Alias  Species	E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester .  Liquid  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified  IgG  Polyclonal  Scavenger receptor class B member 1 (SRB1) (CD36 and LIMPII analogous 1) (CLA-1) (CD36 antigen-like 1) (Collagen type I receptor, thrombospondin receptor-like 1) (SR-BI) (CD antigen CD36), SCARB1, CD36L1 CLA1  Homo sapiens (Human)




## **CUSABIO TECHNOLOGY LLC**











Immunohistochemistry of paraffin-embedded human adrenal gland tissue using CSB-PA845139LA01HU at dilution of 1:100