






GNPNAT1 Antibody

Storage Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Uniprot No. Q96EK6 Immunogen Recombinant Human Glucosamine 6-phosphate N-acetyltransferase protein (1-184AA) Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200 Relevance cytosol, glucosamine 6-phosphate N-acetyltransferase activity, identical protein binding, cellular protein metabolic process, dolichol-linked oligosaccharide biosynthetic process, post-translational protein modification, protein N-linked glycosylation via asparagine, UDP-N-acetylglucosamine biosynthetic process Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4) (Phosphoglucosamine transacetylase),	Product Code	CSB-PA856926LA01HU
Immunogen Recombinant Human Glucosamine 6-phosphate N-acetyltransferase protein (1-184AA) Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200 Relevance cytosol, glucosamine 6-phosphate N-acetyltransferase activity, identical protein binding, cellular protein metabolic process, dolichol-linked oligosaccharide biosynthetic process, post-translational protein modification, protein N-linked glycosylation via asparagine, UDP-N-acetylglucosamine biosynthetic process Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200 Relevance cytosol, glucosamine 6-phosphate N-acetyltransferase activity, identical protein binding, cellular protein metabolic process, dolichol-linked oligosaccharide biosynthetic process, post-translational protein modification, protein N-linked glycosylation via asparagine, UDP-N-acetylglucosamine biosynthetic process Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Uniprot No.	Q96EK6
Species Reactivity Human, Mouse Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200 Relevance cytosol, glucosamine 6-phosphate N-acetyltransferase activity, identical protein binding, cellular protein metabolic process, dolichol-linked oligosaccharide biosynthetic process, post-translational protein modification, protein N-linked glycosylation via asparagine, UDP-N-acetylglucosamine biosynthetic process Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Immunogen	·
Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200 Relevance cytosol, glucosamine 6-phosphate N-acetyltransferase activity, identical protein binding, cellular protein metabolic process, dolichol-linked oligosaccharide biosynthetic process, post-translational protein modification, protein N-linked glycosylation via asparagine, UDP-N-acetylglucosamine biosynthetic process Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method Solution Solutio	Raised In	Rabbit
Relevance cytosol, glucosamine 6-phosphate N-acetyltransferase activity, identical protein binding, cellular protein metabolic process, dolichol-linked oligosaccharide biosynthetic process, post-translational protein modification, protein N-linked glycosylation via asparagine, UDP-N-acetylglucosamine biosynthetic process Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Species Reactivity	Human, Mouse
binding, cellular protein metabolic process, dolichol-linked oligosaccharide biosynthetic process, post-translational protein modification, protein N-linked glycosylation via asparagine, UDP-N-acetylglucosamine biosynthetic process Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200
ConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasGlucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Relevance	binding, cellular protein metabolic process, dolichol-linked oligosaccharide biosynthetic process, post-translational protein modification, protein N-linked
Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Conjugate	Non-conjugated
Isotype IgG Clonality Polyclonal Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Storage Buffer	
Clonality Polyclonal Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Purification Method	>95%, Protein G purified
Alias Glucosamine 6-phosphate N-acetyltransferase (EC 2.3.1.4)	Isotype	IgG
	Clonality	Polyclonal
GNPNAT1, GNA1	Alias	(Phosphoglucosamine acetylase) (Phosphoglucosamine transacetylase),
Species Human	Species	Human
Research Area Others	Research Area	Others
Target Names GNPNAT1	Target Names	GNPNAT1

Image



Western Blot

Positive WB detected in: MCF-7 whole cell lysate, Mouse liver tissue, Mouse kidney tissue,

Mouse stomach tissue

All lanes: GNPNAT1 antibody at 3.4µg/ml


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 21 kDa Observed band size: 21 kDa










Immunohistochemistry of paraffin-embedded human gastric cancer using CSB-PA856926LA01HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human lung cancer using CSB-PA856926LA01HU at dilution of 1:100