





HSD17B14 Antibody


Clonality Polyclonal 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names HSD17B14		
Storage Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Uniprot No. Q9BPX1 Immunogen Recombinant Human 17-beta-hydroxysteroid dehydrogenase 14 protein (1-270AA) Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB; Recommended dilution: WB:1:1000-1:5000 Relevance Has NAD-dependent 17-beta-hydroxysteroid dehydrogenase activity. Converts oestradiol to oestrone. The physiological substrate is not known. Acts on oestradiol and 5-androstene-3-beta,17-beta-diol (in vitro). Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names	Product Code	CSB-PA859026HA01HU
Uniprot No. Q9BPX1 Immunogen Recombinant Human 17-beta-hydroxysteroid dehydrogenase 14 protein (1-270AA) Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB; Recommended dilution: WB:1:1000-1:5000 Relevance Has NAD-dependent 17-beta-hydroxysteroid dehydrogenase activity. Converts oestradiol to oestrone. The physiological substrate is not known. Acts on oestradiol and 5-androstene-3-beta,17-beta-diol (in vitro). Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names	Abbreviation	17-beta-hydroxysteroid dehydrogenase 14
Immunogen Recombinant Human 17-beta-hydroxysteroid dehydrogenase 14 protein (1-270AA) Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB; Recommended dilution: WB:1:1000-1:5000 Relevance Has NAD-dependent 17-beta-hydroxysteroid dehydrogenase activity. Converts oestradiol to oestrone. The physiological substrate is not known. Acts on oestradiol and 5-androstene-3-beta,17-beta-diol (in vitro). Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names HSD17B14	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB; Recommended dilution: WB:1:1000-1:5000 Relevance Has NAD-dependent 17-beta-hydroxysteroid dehydrogenase activity. Converts oestradiol to oestrone. The physiological substrate is not known. Acts on oestradiol and 5-androstene-3-beta,17-beta-diol (in vitro). Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names HSD17B14	Uniprot No.	Q9BPX1
Species Reactivity Human, Mouse Tested Applications ELISA, WB; Recommended dilution: WB:1:1000-1:5000 Relevance Has NAD-dependent 17-beta-hydroxysteroid dehydrogenase activity. Converts oestradiol to oestrone. The physiological substrate is not known. Acts on oestradiol and 5-androstene-3-beta,17-beta-diol (in vitro). Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names HSD17B14	Immunogen	
Tested Applications ELISA, WB; Recommended dilution: WB:1:1000-1:5000 Relevance Has NAD-dependent 17-beta-hydroxysteroid dehydrogenase activity. Converts oestradiol to oestrone. The physiological substrate is not known. Acts on oestradiol and 5-androstene-3-beta,17-beta-diol (in vitro). Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names	Raised In	Rabbit
Relevance Has NAD-dependent 17-beta-hydroxysteroid dehydrogenase activity. Converts oestradiol to oestrone. The physiological substrate is not known. Acts on oestradiol and 5-androstene-3-beta,17-beta-diol (in vitro). Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300	Species Reactivity	Human, Mouse
oestradiol to oestrone. The physiological substrate is not known. Acts on oestradiol and 5-androstene-3-beta,17-beta-diol (in vitro). Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names HSD17B14	Tested Applications	ELISA, WB; Recommended dilution: WB:1:1000-1:5000
ConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAlias17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1SpeciesHumanResearch AreaSignal TransductionTarget NamesHSD17B14	Relevance	oestradiol to oestrone. The physiological substrate is not known. Acts on
Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names HSD17B14	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names HSD17B14	Conjugate	Non-conjugated
Isotype IgG Clonality Polyclonal Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names HSD17B14	Storage Buffer	
Clonality Polyclonal 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names HSD17B14	Purification Method	>95%, Protein G purified
Alias 17-beta-hydroxysteroid dehydrogenase 14 (17-beta-HSD 14) (EC 1.1.1) (17-beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction HSD17B14	Isotype	IgG
beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14, DHRS10 SDR3 SDR47C1 Species Human Research Area Signal Transduction Target Names HSD17B14	Clonality	Polyclonal
Research Area Signal Transduction Target Names HSD17B14	Alias	beta-hydroxysteroid dehydrogenase DHRS10) (Dehydrogenase/reductase SDR family member 10) (Retinal short-chain dehydrogenase/reductase retSDR3) (Short chain dehydrogenase/reductase family 47C member 1), HSD17B14,
Target Names HSD17B14	Species	Human
	Research Area	Signal Transduction
Income.	Target Names	HSD17B14
ımage	Image	



CUSABIO TECHNOLOGY LLC







Western blot

All lanes: HSD17B14 antibody at 2µg/ml +

Mouse kidney tissue

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 29 kDa Observed band size: 29 kDa