



HSD17B10 Antibody, FITC conjugated

Product Code	CSB-PA860776HC01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q99714
Immunogen	Recombinant Human 3-hydroxyacyl-CoA dehydrogenase type-2 protein (2-261AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	Functions in mitochondrial tRNA maturation. Part of mitochondrial ribonuclease P, an enzyme composed of MRPP1/TRMT10C, MRPP2/HSD17B10 and MRPP3/KIAA0391, which cleaves tRNA molecules in their 5'-ends. Catalyzes the beta-oxidation at position 17 of androgens and estrogens and has 3-alpha-hydroxysteroid dehydrogenase activity with androsterone. Catalyzes the third step in the beta-oxidation of fatty acids. Carries out oxidative conversions of 7-alpha-OH and 7-beta-OH bile acids. Also exhibits 20-beta-OH and 21-OH dehydrogenase activities with C21 steroids. By interacting with intracellular amyloid-beta, it may contribute to the neuronal dysfunction associated with Alzheimer disease (AD).
Form	Liquid
Conjugate	FITC
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	3-hydroxyacyl-CoA dehydrogenase type-2 (EC 1.1.1.35) (17-beta-hydroxysteroid dehydrogenase 10) (17-beta-HSD 10) (EC 1.1.1.51) (2-methyl-3-hydroxybutyryl-CoA dehydrogenase) (MHBD) (3-hydroxy-2-methylbutyryl-CoA dehydrogenase) (EC 1.1.1.178) (3-hydroxyacyl-CoA dehydrogenase type II) (Endoplasmic reticulum-associated amyloid beta-peptide-binding protein) (Mitochondrial ribonuclease P protein 2) (Mitochondrial RNase P protein 2) (Short chain dehydrogenase/reductase family 5C member 1) (Short-chain type dehydrogenase/reductase XH98G2) (Type II HADH), HSD17B10, ERAB HADH2 MRPP2 SCHAD SDR5C1 XH98G2
Species	Homo sapiens (Human)
Research Area	Neuroscience
Target Names	HSD17B10