






KCNK13 Antibody

StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.Q9HB14ImmunogenRecombinant Human Potassium channel subfamily K member 13 protein (292-408AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200, IF:1:50-1:200RelevancePotassium channel displaying weak inward rectification in symmetrical K(+) solution.FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPotassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13SpeciesHumanResearch AreaNeuroscienceTarget NamesKCNK13	Product Code	CSB-PA862058LA01HU
ImmunogenRecombinant Human Potassium channel subfamily K member 13 protein (292-408AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200, IF:1:50-1:200RelevancePotassium channel displaying weak inward rectification in symmetrical K(+) solution.FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPotassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13SpeciesHumanResearch AreaNeuroscience	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised In Rabbit Species Reactivity Human Tested Applications ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:2000, IF:1:50-1:2000 Relevance Potassium channel displaying weak inward rectification in symmetrical K(+) solution. Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Potassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13 Species Human Research Area Neuroscience	Uniprot No.	Q9HB14
Species ReactivityHumanTested ApplicationsELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200, IF:1:50-1:200RelevancePotassium channel displaying weak inward rectification in symmetrical K(+) solution.FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPotassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13SpeciesHumanResearch AreaNeuroscience	Immunogen	, i
Tested ApplicationsELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200RelevancePotassium channel displaying weak inward rectification in symmetrical K(+) solution.FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPotassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13SpeciesHumanResearch AreaNeuroscience	Raised In	Rabbit
Relevance Potassium channel displaying weak inward rectification in symmetrical K(+) solution. Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Potassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13 Species Human Research Area Neuroscience	Species Reactivity	Human
Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Potassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13 Species Human Research Area Neuroscience	Tested Applications	
ConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasPotassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13SpeciesHumanResearch AreaNeuroscience	Relevance	
Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Potassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13 Species Human Research Area Neuroscience	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Potassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13 Species Human Research Area Neuroscience	Conjugate	Non-conjugated
Isotype IgG Clonality Polyclonal Alias Potassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13 Species Human Research Area Neuroscience	Storage Buffer	
Clonality Polyclonal Alias Potassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13 Species Human Research Area Neuroscience	Purification Method	>95%, Protein G purified
Alias Potassium channel subfamily K member 13 (Tandem pore domain halothane-inhibited potassium channel 1) (THIK-1), KCNK13 Species Human Neuroscience	Isotype	IgG
inhibited potassium channel 1) (THIK-1), KCNK13 Species Human Research Area Neuroscience	Clonality	Polyclonal
Research Area Neuroscience	Alias	· · · · · · · · · · · · · · · · · · ·
	Species	Human
Target Names KCNK13	Research Area	Neuroscience
	Target Names	KCNK13





Western blot

All lanes: KCNK13 antibody at 8µg/ml Lane 1: HepG2 whole cell lysate Lane 2: LO2 whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 46 kDa Observed band size: 46 kDa
















Immunohistochemistry of paraffin-embedded human skeletal muscle tissue using CSB-PA862058LA01HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human glioma using CSB-PA862058LA01HU at dilution of 1:100



Immunofluorescent analysis of HepG2 cells using CSB-PA862058LA01HU at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)