



MED28 Antibody, FITC conjugated

Product Code	CSB-PA863955LC01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9H204
Immunogen	Recombinant Human Mediator of RNA polymerase II transcription subunit 28 protein (1-178AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton downstream of tyrosine kinase signaling pathways.
Form	Liquid
Conjugate	FITC
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Mediator of RNA polymerase II transcription subunit 28 (Endothelial-derived protein 1) (Mediator complex subunit 28) (Merlin and Grb2-interacting cytoskeletal protein) (Magicin) (Tumor angiogenesis marker EG-1), MED28, EG1
Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	MED28