





UBE2O Antibody


Product Code	CSB-PA866339LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9C0C9
Immunogen	Recombinant Human (E3-independent) E2 ubiquitin-conjugating enzyme protein (10-277AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB; Recommended dilution: WB:1:500-1:2000
Relevance	E2/E3 hybrid ubiquitin-protein ligase that displays both E2 and E3 ligase activities and mediates monoubiquitination of target proteins (PubMed:23455153, PubMed:24703950). Negatively regulates TRAF6-mediated NF-kappa-B activation independently of its E2 activity (PubMed:23381138). Acts as a positive regulator of BMP7 signaling by mediating monoubiquitination of SMAD6, thereby regulating adipogenesis (PubMed:23455153). Mediates monoubiquitination at different sites of the nuclear localization signal (NLS) of BAP1, leading to cytoplasmic retention of BAP1. Also able to monoubiquitinate the NLS of other chromatin-associated proteins, such as INO80 and CXXC1, affecting their subcellular location (PubMed:24703950). Acts as a regulator of retrograde transport by assisting the TRIM27:MAGEL2 E3 ubiquitin ligase complex to mediate \\\'Lys-63\\\'-linked ubiquitination of WASH1, leading to promote endosomal F-actin assembly (PubMed:23452853).
Form	Liquid
Form Conjugate	Liquid Non-conjugated
	•
Conjugate	Non-conjugated Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Antigen Affinity Purified
Conjugate Storage Buffer Purification Method Isotype	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Antigen Affinity Purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	Non-conjugated Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Antigen Affinity Purified IgG Polyclonal (E3-independent) E2 ubiquitin-conjugating enzyme (EC 2.3.2.24) (E2/E3 hybrid ubiquitin-protein ligase UBE2O) (Ubiquitin carrier protein O) (Ubiquitin-conjugating enzyme E2 of 230 kDa) (Ubiquitin-conjugating enzyme E2-230K) (Ubiquitin-protein ligase O), UBE2O,
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Antigen Affinity Purified IgG Polyclonal (E3-independent) E2 ubiquitin-conjugating enzyme (EC 2.3.2.24) (E2/E3 hybrid ubiquitin-protein ligase UBE2O) (Ubiquitin carrier protein O) (Ubiquitin-conjugating enzyme E2 of 230 kDa) (Ubiquitin-conjugating enzyme E2-230K) (Ubiquitin-protein ligase O), UBE2O, KIAA1734
Conjugate Storage Buffer Purification Method Isotype Clonality Alias Species	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Antigen Affinity Purified IgG Polyclonal (E3-independent) E2 ubiquitin-conjugating enzyme (EC 2.3.2.24) (E2/E3 hybrid ubiquitin-protein ligase UBE2O) (Ubiquitin carrier protein O) (Ubiquitin-conjugating enzyme E2 of 230 kDa) (Ubiquitin-conjugating enzyme E2 of 230 kDa) (Ubiquitin-conjugating enzyme E2-230K) (Ubiquitin-protein ligase O), UBE2O, KIAA1734 Human



CUSABIO TECHNOLOGY LLC







Western blot

All lanes: UBE2O antibody at 0.8µg/ml + A431

whole cell lysate Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 142 kDa Observed band size: 142 kDa