





SMURF2 Antibody, Biotin conjugated

Product Code	CSB-PA872532ED01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9HAU4
Immunogen	Recombinant Human E3 ubiquitin-protein ligase SMURF2 protein (1-748AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Interacts with SMAD1 and SMAD7 in order to trigger their ubiquitination and proteasome-dependent degradation. In addition, interaction with SMAD7 activates autocatalytic degradation, which is prevented by interaction with SCYE1. Forms a stable complex with the TGF-beta receptor-mediated phosphorylated SMAD2 and SMAD3. In this way, SMAD2 may recruit substrates, such as SNON, for ubiquitin-mediated degradation. Enhances the inhibitory activity of SMAD7 and reduces the transcriptional activity of SMAD2. Coexpression of SMURF2 with SMAD1 results in considerable decrease in steady-state level of SMAD1 protein and a smaller decrease of SMAD2 level.
Form	Liquid
Form Conjugate	· · · · · · · · · · · · · · · · · · ·
	Liquid
Conjugate	Liquid Biotin Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Conjugate Storage Buffer Purification Method Isotype	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal E3 ubiquitin-protein ligase SMURF2 (hSMURF2) (EC 2.3.2.26) (HECT-type E3 ubiquitin transferase SMURF2) (SMAD ubiquitination regulatory factor 2)
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal E3 ubiquitin-protein ligase SMURF2 (hSMURF2) (EC 2.3.2.26) (HECT-type E3 ubiquitin transferase SMURF2) (SMAD ubiquitination regulatory factor 2) (SMAD-specific E3 ubiquitin-protein ligase 2), SMURF2