






## SLC6A15 Antibody

Product Code	CSB-PA875657LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9H2J7
Immunogen	Recombinant Human Sodium-dependent neutral amino acid transporter B(0)AT2 protein (357-452AA)
Raised In	Rabbit
Species Reactivity	Human
<b>Tested Applications</b>	ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200
Relevance	Functions as a sodium-dependent neutral amino acid transporter. Exhibits preference for the branched-chain amino acids, particularly leucine, valine and isoleucine and methionine. Mediates the saturable, pH-sensitive and electrogenic cotransport of proline and sodium ions with a stoichiometry of 1:1. May have a role as transporter for neurotransmitter precursors into neurons. In contrast to other members of the neurotransmitter transporter family, does not appear to be chloride-dependent.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Sodium-dependent neutral amino acid transporter B(0)AT2 (Sodium- and chloride-dependent neurotransmitter transporter NTT73) (Sodium-coupled branched-chain amino-acid transporter 1) (Solute carrier family 6 member 15) (Transporter v7-3), SLC6A15, B0AT2 NTT73 SBAT1
Species	Human
Research Area	Signal Transduction
Target Names	SLC6A15
Image	Immunohistochemistry of paraffin-embedded



Immunohistochemistry of paraffin-embedded human bladder cancer using CSB-PA875657LA01HU at dilution of 1:100















Immunohistochemistry of paraffin-embedded human prostate cancer using CSB-PA875657LA01HU at dilution of 1:100



Immunofluorescent analysis of HepG2 cells using CSB-PA875657LA01HU at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)