






XPNPEP1 Antibody

StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.Q9NQW7ImmunogenRecombinant Human Xaa-Pro aminopeptidase 1 protein (2-623AA)Raised InRabbitSpecies ReactivityHuman, MouseTested ApplicationsELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200RelevanceContributes to the degradation of bradykinin. Catalyzes the removal of a penultimate prolyl residue from the N-termini of peptides, such as Arg-Pro-Pro.FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasXaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase P) (X-Prolyl aminopeptidase 1, soluble), XPNPEP1, XPNPEP	Product Code	CSB-PA878859HA01HU
ImmunogenRecombinant Human Xaa-Pro aminopeptidase 1 protein (2-623AA)Raised InRabbitSpecies ReactivityHuman, MouseTested ApplicationsELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200RelevanceContributes to the degradation of bradykinin. Catalyzes the removal of a penultimate prolyl residue from the N-termini of peptides, such as Arg-Pro-Pro.FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasXaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEP1, XPNPEPL XPNPEPL XPNPEPL XPNPEPL APNPEPL	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitSpecies ReactivityHuman, MouseTested ApplicationsELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200RelevanceContributes to the degradation of bradykinin. Catalyzes the removal of a penultimate prolyl residue from the N-termini of peptides, such as Arg-Pro-Pro.FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasXaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEP1, XPNPEPL XPNPEPL1SpeciesHumanResearch AreaSignal Transduction	Uniprot No.	Q9NQW7
Species ReactivityHuman, MouseTested ApplicationsELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200RelevanceContributes to the degradation of bradykinin. Catalyzes the removal of a penultimate prolyl residue from the N-termini of peptides, such as Arg-Pro-Pro.FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasXaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (SAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEP1, XPNPEPL XPNPEPL1SpeciesHumanResearch AreaSignal Transduction	Immunogen	Recombinant Human Xaa-Pro aminopeptidase 1 protein (2-623AA)
Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200 Relevance Contributes to the degradation of bradykinin. Catalyzes the removal of a penultimate prolyl residue from the N-termini of peptides, such as Arg-Pro-Pro. Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Xaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEP1, XPNPEPL XPNPEPL1 Species Human Research Area Signal Transduction	Raised In	Rabbit
Relevance Contributes to the degradation of bradykinin. Catalyzes the removal of a penultimate prolyl residue from the N-termini of peptides, such as Arg-Pro-Pro. Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Xaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEPL, XPNPEPL1 Species Human Research Area Signal Transduction	Species Reactivity	Human, Mouse
penultimate prolyl residue from the N-termini of peptides, such as Arg-Pro-Pro. Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300	Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200
ConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasXaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEP1, XPNPEPL XPNPEPL1SpeciesHumanResearch AreaSignal Transduction	Relevance	·
Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Xaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEP1, XPNPEPL XPNPEPL1 Species Human Research Area Signal Transduction	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Isotype IgG Clonality Polyclonal Alias Xaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEPL, XPNPEPL1 Species Human Research Area Signal Transduction	Conjugate	Non-conjugated
Isotype IgG Clonality Polyclonal Alias Xaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEP1, XPNPEPL XPNPEPL1 Species Human Research Area Signal Transduction	Storage Buffer	
Clonality Polyclonal Alias Xaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEPL, XPNPEPL XPNPEPL1 Species Human Research Area Signal Transduction	Purification Method	>95%, Protein G purified
Alias Xaa-Pro aminopeptidase 1 (EC 3.4.11.9) (Aminoacylproline aminopeptidase) (Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEP1, XPNPEPL XPNPEPL1 Species Human Research Area Signal Transduction	Isotype	IgG
(Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro aminopeptidase 1) (X-prolyl aminopeptidase 1, soluble), XPNPEP1, XPNPEPL XPNPEPL1 Species Human Research Area Signal Transduction	Clonality	Polyclonal
Research Area Signal Transduction	Alias	(Cytosolic aminopeptidase P) (Soluble aminopeptidase P) (sAmp) (X-Pro
		XPNPEPL1
Target Names XPNPEP1	Species	
	-	Human





Western blot

All lanes: Xaa-Pro aminopeptidase 1 antibody at

8µg/ml

Lane 1: Mouse small intestine tissue Lane 2: Mouse stomach tissue Lane 3: Mouse kidney tissue

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 70, 68, 75, 73 kDa

Observed band size: 70 kDa




CUSABIO TECHNOLOGY LLC











Immunohistochemistry of paraffin-embedded human colon cancer using CSB-PA878859HA01HU at dilution of 1:100