





SMPD3 Antibody, Biotin conjugated

Product Code	CSB-PA878920LD01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9NY59
Immunogen	Recombinant Human Sphingomyelin phosphodiesterase 3 protein (401-655AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA
Relevance	Catalyzes the hydrolysis of sphingomyelin to form ceramide and phosphocholine. Ceramide mediates numerous cellular functions, such as apoptosis and growth arrest, and is capable of regulating these 2 cellular events independently. Also hydrolyzes sphingosylphosphocholine. Regulates the cell cycle by acting as a growth suppressor in confluent cells. Probably acts as a regulator of postnatal development and participates in bone and dentin
	mineralization.
Form	mineralization. Liquid
Form Conjugate	
	Liquid
Conjugate	Liquid Biotin Preservative: 0.03% Proclin 300
Conjugate Storage Buffer	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Conjugate Storage Buffer Purification Method	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Conjugate Storage Buffer Purification Method Isotype	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG
Conjugate Storage Buffer Purification Method Isotype Clonality	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Sphingomyelin phosphodiesterase 3 (EC 3.1.4.12) (Neutral sphingomyelinase
Conjugate Storage Buffer Purification Method Isotype Clonality Alias	Liquid Biotin Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Sphingomyelin phosphodiesterase 3 (EC 3.1.4.12) (Neutral sphingomyelinase 2) (nSMase-2) (nSMase-2) (Neutral sphingomyelinase II), SMPD3