




TNFRSF10D Antibody

Product Code	CSB-PA887006ESR1HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9UBN6
Immunogen	Recombinant Human Tumor necrosis factor receptor superfamily member 10D protein (257-386AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:200-1:1000, IHC:1:20-1:200
Relevance	Receptor for the cytotoxic ligand TRAIL. Contains a truncated death domain and hence is not capable of inducing apoptosis but protects against TRAIL-mediated apoptosis. Reports are contradictory with regards to its ability to induce the NF-kappa-B pathway. According to PubMed:9382840, it cannot but according to PubMed:9430226, it can induce the NF-kappa-B pathway.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	Tumor necrosis factor receptor superfamily member 10D (Decoy receptor 2) (DcR2) (TNF-related apoptosis-inducing ligand receptor 4) (TRAIL receptor 4) (TRAIL-R4) (TRAIL receptor with a truncated death domain) (CD antigen CD264), TNFRSF10D, DCR2 TRAILR4 TRUNDD
Species	Human
Research Area	Cell Biology
Target Names	TNFRSF10D

Image




Western blot


All lanes: Tumor necrosis factor receptor superfamily member 10D antibody at 2µg/ml + Mouse liver tissue

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution
Predicted band size: 42 kDa
Observed band size: 42 kDa



Immunohistochemistry of paraffin-embedded human tonsil tissue using CSB-PA887006ESR1HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human kidney tissue using CSB-PA887006ESR1HU at dilution of 1:100