




FZR1 Antibody

Product Code	CSB-PA892473LA01HU
Abbreviation	Fizzy-related protein homolog
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9UM11
Immunogen	Recombinant Human Fizzy-related protein homolog protein (1-493AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200
Relevance	Key regulator of ligase activity of the anaphase promoting complex/cyclosome (APC/C), which confers substrate specificity upon the complex. Associates with the APC/C in late mitosis, in replacement of CDC20, and activates the APC/C during anaphase and telophase. The APC/C remains active in degrading substrates to ensure that positive regulators of the cell cycle do not accumulate prematurely. At the G1/S transition FZR1 is phosphorylated, leading to its dissociation from the APC/C. Following DNA damage, it is required for the G2 DNA damage checkpoint: its dephosphorylation and reassociation with the APC/C leads to the ubiquitination of PLK1, preventing entry into mitosis.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Fizzy-related protein homolog (Fzr) (CDC20-like protein 1) (Cdh1/Hct1 homolog) (hCDH1), FZR1, CDH1 FYR FZR KIAA1242
Species	Human
Research Area	Cell Biology
Target Names	FZR1
Image	



Western blot

All lanes: FZR1 antibody at 6 μ g/ml

Lane 1: Hela whole cell lysate

Lane 2: Jurkat whole cell lysate


Lane 3: 293T whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 56, 55, 45 kDa

Observed band size: 56 kDa



Immunohistochemistry of paraffin-embedded

human testis tissue using CSB-

PA892473LA01HU at dilution of 1:100