


Image






BLK Antibody

| Product Code | CSB-PA926374 |
|----------------------------|---|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P51451 |
| Immunogen | Fusion protein of Human BLK |
| Raised In | Rabbit |
| Species Reactivity | Human, Mouse |
| Tested Applications | ELISA,WB,IHC;ELISA:1:1000-1:2000,WB:1:200-1:1000,IHC:1:25-1:100 |
| Relevance | This gene encodes a nonreceptor tyrosine-kinase of the src family of proto- oncogenes that are typically involved in cell proliferation and differentiation. The protein has a role in B-cell receptor signaling and B-cell development. The protein also stimulates insulin synthesis and secretion in response to glucose and enhances the expression of several pancreatic beta-cell transcription factors. |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol |
| Purification Method | Antigen affinity purification |
| Isotype | IgG |
| Species | Homo sapiens (Human) |
| Target Names | BLK |
| _ | |



The image on the left is immunohistochemistry of paraffin-embedded Human gastic cancer tissue using CSB-PA926374(BLK Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: x200)



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using CSB-PA926374(BLK Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: x200)




CUSABIO TECHNOLOGY LLC









Gel: 8%SDS-PAGE, Lysate: 40 µg, Lane: Raji cells, Primary antibody: CSB-PA926374(BLK Antibody) at dilution 1/250, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 20 seconds